

Europäisches Patentamt

European Patent Office

Office européen des brevets

-15 04 2004

Best Available Copy

REC'D 0 3 MAY 2004
WIPO PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent

Patent application No. Demande de brevet nº

03005181.7

Der Präsident des Europäischen Patentamts; Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets p.o.

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN COMPLIANCE WITH RULE 17.1(a) OR (b)

R C van Dijk



Anmeldung Nr:

Application no.:

03005181.7

Demande no:

Anmeldetag:

Date of filing:

07.03.03

Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

BASF Plant Science GmbH

67056 Ludwigshafen ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention: (Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung. If no title is shown please refer to the description. Si aucun titre n'est indiqué se referer à la description.)

Enhanced amylose production in plants

In Anspruch genommene Prioriät(en) / Priority(ies) claimed /Priorité(s) revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT SE SI SK TR LI

Enhanced amylose production in plants

Description

The present invention relates to new amylose biosynthesis enhancing proteins, nucleic acids encoding a starch biosynthesis enhancing protein, a method for producing amylose with high efficiency by culturing genetically modified plants with an increased amylose biosynthesis compared to the wild type or to the genetically modified plants themselves as well as the use of these transgenic plants over-expressing at least one of the amylose biosynthesis enhancing proteins for the production of amylose.

Starch is the major storage carbohydrate of plants and is mainly accumulated in seeds and tubers, which are then the reproductive tissues of plants that form those types of organs. Starch is also accumulated on a diurnal basis where starch is built up in green tissue from photosynthetic products and then metabolised for energy during the dark period. The storage starch is assembled into semi crystalline granules. Amylopectin and amylose are the two constituent molecules of starch. Amylopectin is a branched molecule consisting of linear α -1,4 glucan chains linked by α -1,6 bonds. Amylose consists essentially of the linear α -1,4 glucan chains.

20

25

30

15

Starch is utilised for many applications within the technical industry as well as the food industry. Main crops used by starch processors are maize and potato. For potatoes specific varieties are utilised for starch production that have been bred for high starch contents. This means that the starch content and yield is an important economic driver for the starch processing industry. A greater part of produced dry starch is used for paper production. The specifications and requirements for the starch component varies from application to application and starch is many times chemically modified in order to provide desired properties to an application. Another way to achieve starch of different qualities is to take advantage of mutations in the starch biosynthesis and more recently by genetic modification of pathways leading to starch. The first main modifications have been to separate the production of the two starch components amylopectin and amylose into different varieties. Waxy or "amylose free" varieties contain solely amylopectin type starch while there are also high amylose genotypes such as "amylose extender" in maize.

35

40

Amylose starch has several potential industrial uses as a film former or for expanded products. High amylose starch can be achieved in potatoes and other starch containing plants by inhibition of starch branching enzymes. This leads then to the concomitant reduction or elimination of amylopectin branching and thereby an increased amylose

10

15

20

25

30

35

40

US 5,856,467 describes the genetically engineered modification of potato for suppressing formation of amylopectin-type starch. The document describes an antisense construct for inhibiting, to a varying extent, the expression of the gene coding for formation of starch branching enzyme (SBE gene) in potato, said antisense construct comprising a tuber specific promoter, transcription start and the first exon of the SBE gene, inserted in the antisense direction.

US 6,169,226 relates to an amino acid sequence of a second starch branching enzyme (SBE II) of potato and a fragment thereof as well as to the corresponding isolated DNA sequences. It describes the production of transgenic potatoes and the use of these transgenic potatoes for the production of amylose-type starch.

WO 97/20040 and WO 98/20145 describe methods of altering the amylopectine/ amylose starch content of plant cells by introducing into the plant cells nucleic acid sequences operably linked in sense or antisense orientation to a suitable promoter which homologous genes encodes polypeptides having SBE I or SBE II activity.

A side effect of the amylose overproduction is a decreased total starch content in the potatoes. This decrease becomes more pronounced as the amylose fraction is increased.

Basic enzymes for the production of amylopectin and amylose are starch synthases that build the linear α -1,4 glucan chains and branching enzymes breaking the α -1,4 glucan chain and reattaching them by α -1,6 bonds. Several other enzymes are likely to affect starch structure and composition, such as debranching enzymes, but initially most focus has been towards affecting the expression of starch synthases and starch branching enzymes. This has led to an extensive dissection of what enzymes are important for what features of starch synthesis. However it has never been convincingly shown how the synthesis of starch in plants whether amylose or amylopectin is initiated.

Suggestions on the initiation of starch biosynthesis have been the subject of several scientific papers since it has been difficult to attribute a primer independent function to starch synthases under other than artificial in vitro conditions. By primer independent function implies the formation of new α -1,4 glucan chains with ADP-glucose as the sole starting point and building block. One proposed pathway has been that the presence of maltooligosaccharides act as primers for the addition of further glucose units by starch synthases although it has been debated on whether concentrations are sufficient to provide the basis for starch synthesis and also how these maltooligosaccharides would be formed in the plastids.

Starch is in plants synthesised as an energy storage molecule. Much is known about the enzymes participating in the starch biosynthesis although, the initiation of the starch molecule. In mammalians and yeast an energy storage molecule very similar to starch is synthesised, glycogen. The enzymatic steps for synthesis of the respective molecules are analogous. In glycogen biosynthesis the initiation of the molecule is known and synthesised by the enzyme glycogenin. Glycogenin is a self-glucosylating enzyme polymerising a linear chain of approximately 8 glucose molecules on itself. The primer of about 8 glucose residues is necessary for the enzymes catalysing the continuation of glucose incorporation to the glycogen molecule to function.

Cheng et al., 1995, Mol. and Cell. Biol. 6632-6640 compare the two yeast proteins with rabbit muscle glycogenin.

15 Roach et al., 1997, Progress in Nucleic Acid Research and Molecular Biology Vol 57, describe self glycosylating initiator proteins and their roll in glycogen biosynthesis.

Mu et al., 1997, Journal of Biological Chemistry 272 (44), 27589-27597 compare mammalian with yeast and C. elegans glycogenins.

Factors important for starch quantity have been investigated and many initiatives have been taken, especially in potato, to Increase starch formation and content by over-expression or inhibition of various enzyme activities in areas of increased substrate supply, increased biosynthesis activity or shutting down substrate diverting pathways but so far this has led only to limited success with no commercial applications and only some scientific publications.

Regierer, B. et al., Starch content and yield increase as a result of altering adenylate pools in transgenic plants. Nat Biotechnol. 20(12):1256-60, (2002).

Sweetlove, LJ et al., Starch synthesis in transgenic potato tubers with increased 3-phosphoglyceric acid content as a consequence of increased 6-phosphofructokinase activity. Planta 213(3):478-82 (2001).

Veramendi, J et al.., Antisense repression of hexokinase 1 leads to an overaccumulation of starch in leaves of transgenic potato plants but not to significant changes in tuber carbohydrate metabolism. Plant Physiol. 121(1):123-34 (1999).

Geigenberger, P et al.., Overexpression of pyrophosphatase leads to increased sucrose degradation and starch synthesis, increased activities of enzymes for sucrose-

10

20

25

10

15

20

25

30

starch interconversions, and increased levels of nucleotides in growing potato tubers. Planta. 205(3):428-37(1998).

Sweetlove, LJ et al., Starch metabolism in tubers of transgenic potato (Solanum tuberosum) with increased ADPglucose pyrophosphorylase. Biochem J. 320 (Pt 2):493-8 (1996).

In other research a biochemical function superficially similar to the one initiating glycogen production in animals was investigated. A class of genes have then been isolated from several plants and was given the name amylogenin (WO94/04693; Sing, D. et al, β-Glucosylarginin: a new glucose-protein bond in a self-glucosylating protein from sweet corn, FEBS Letters 376:61-64, (1995) in the belief that it was the plant equivalent of glycogenin which acts as a self-glycosylating enzyme and provide primers for starch biosynthesis in plants. These genes have no resemblance from a structural point of view to the genes coding for glycogenin and have later been determined not to have a function in starch biosynthesis but rather might be of importance for cell wall formation , see Bocca, S.N et al., Molecular cloning and characterization of the enzyme UDP-glucose: protein transglucosylase from potato. Plant Physiology and Biochemistry 37(11):809-819(1999).

WO 98/50553 describes nucleic acid fragments encoding a plant glycogenin or a water stress protein. WO 98/50553 also relates to the construction of chimeric genes encoding all or a portion of a plant glycogenin in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of a plant glycogenin in a transformed host cell.

Thus although many enzymes and pathways have been investigated in plants, the question on how starch formation is initiated and what determines the starch content is still unresolved.

Amylose is a commercially important starch product with many uses but unfortunately an increase in amylose content in transgenic potato plants is associated with a significant decrease in starch content, see figure 1.

Analyses of transgenic high amylose potato lines show that there is an excess of soluble sugars in these lines, see figure 2. This indicates that the starch biosynthesis in these transgenic lines is not efficient enough for incorporation of available sugars.

Amylose starch consists of very few reducing ends compared to native starch. There-40 fore it is commercially important to identify genes that further enhance the amylose

1 F - OT COOP_MAIL_10

20

30

35

biosynthesis and that are capable to incorporate the excess of glucose residues available and to compensate the decrease in starch content in plants that produces amylose in high amounts.

The invention aims at enhancing the yield of amylose biosynthesis by the overexpression of genes which enhance amylose biosynthesis in transformed plants.

The invention describes genes coding for proteins which enhance amylose production.

The present invention describes the nucleic acids SEQ ID NO 1 and 3 from potato coding for enzymes enhancing the de novo amylose biosynthesis.

Example 1 describes that the nucleic acid sequences SEQ ID NO 1 or 3 can complement a missing glycogenin function in yeast cells containing knock-out mutations for the self-glycosylating proteins Glg1p and Glg2p.

Gene constructs were made for gene-inhibition and over-expression of the two genes SEQ ID NO 1 or 3 in potato. Transgenic lines with the over-expressed or inhibited enzyme activity were analysed with regard to the genes influence on amylose content.

Both genes were inserted in sense and antisense direction downstream of a plant promoter element, resulting in the transformation binary vectors pHS1, pHS2, pHS3 and pHS4, see figures 3-7.

25 The antisense constructs were transformed into potato plant varieties Prevalent and Producent and the sense constructs were transformed to the potato varieties Desiree and the transgenic plant AM99-2003 according to the transformation method as described in example 2. The transgenic plant AM99-2003 was produced as described in example 3.

Prevalent and Producent are starch varieties having a starch content of approximately 20 %. Desirée is a potato variety having a starch content of approximately 16% and AM99-2003 is a transgenic high amylose line having a starch and thereby amylose content of approximately 13%.

The putative genes were Isolated from a tuber specific cDNA library of Solanum tuberosum (variety Prevalent). The library was made from a lambdaZAP directional kit (Stratagene).

Both cDNAs isolated were full-length clones of the individual genes and named StGH1 and StGH2, for nucleic acid sequences see SEQ ID NO 1 and SEQ ID NO 3.

pHS1

5

10

20 ·

A 1300bp PCR fragment from the StGH1 gene was constructed in antisense direction driven by the gbss promoter. The PCR fragment was cut out from its cloning vector pCR4-TOPO (Invitrogen) with EcoRI (blunted) and Xbal. The fragment was ligated to the pGPTV-kan (Becker, D. et al., Plant Molecular Biology 20:1195-1197(1992) based binary vector pHo3.1 between a gbss promoter (WO 92/11376) and a nos terminator at the Sall (blunted) and Xbal sites. The binary vector also includes nptII as selection marker driven by the nos promoter (Herrera, L. et al., 1983). The construct was named pHS1, for details see figure 3a and 4.

15 pHS2

A 2300bp full-length cDNA clone of StGH2 was cut out from the cloning vector pBluescript (Stratagene) with Xbal and Xhol. The gene was ligated in antisense direction between the gbss promoter and nos terminator to the binary vector pHo3.1 at Xbal and Sall. As can be seen under pHS1 the vector has nptll as selection system. The vector was named pHS2, for details see figure 3b and 5.

pHS3

- A full-length StGH1 cDNA, (1780bp) was cut out from the host vector pBluescript with EcoRI (blunted) and BgIII and ligated to the BamHI and Smal sites of pUCgbssprom (3886bp), containing pUC19 with the gbss promoter and the nos terminator. The plasmid was named pUCGH1.
- A fragment with the gbss promoter, the StGH1 gene and the nos terminator was moved from pUCGH1 with EcoRI (blunted) and HindIII (2980bp) and ligated to PstI (blunted) and HindIII opened pSUN1 (WO 02/00900). The plasmid was named pSUNGH1.
- A 3600bp fragment containing the AHAS resistance gene from Arabidopsis thaliana (Sathasivan, K. et al., Plant Physiology 97(1991), 1044-1050) with nos promoter, see Herrera-Estrella, L. et al., Nature 303:209-213(1983) and OCS terminator (Wesley, S.V. et al., Plant J. 27(6):581-590(2001) was ligated to pSUNGH1 (9000bp) at the Smal site. The vector was given the name pHS3, for details see figure 3c and 6.

pHS4

The gbss promoter and nos terminator was ligated to pBR322 with EcoRI and HindIII. Between the promoter and terminator an EcoRi-HincII full-length gene pStGH2 was cloned at the Xbal site. The 3366bp promotor-gene-terminator complex was cut using EcoRI (partial digestion) and EcoRV, and ligated to pSUN1 at EcoRi-EcoRV and named pSUNGH2. An Xbal fragment with AHAS gene (Arabidopsis thaliana), nos promoter and OCS terminator was ligated to pSUNGH2 opened with Xbal (partial digestion). The AHAS gene is used as selection marker. The construct was named pHS4, for details see figure 3d and 7.

Example 2 describes the general method for the transformation of different potato plant varieties producing native starch or high amylose type starch with pHS1, pHS2, pHS3 or pHS4.

15

10

The StGH1 and StGH2 genes were down-regulated in the potato plant varieties Prevalent and Producent by transformation with the genes in antisense direction in relation to a plant regulatory element as described in example 4 and 6. Down-regulation of the two genes resulted in a decrease in gene expression in transgenic lines, compared to their mother varieties in the order of 50-95%, see example 7 and table 3. Transgenic lines transformed with pHS1 and pHS2 with confirmed decrease in gene-expression have a decrease in dry matter of 7 to 11% compared to their mother varieties, see example 8 and table 5.

The StGH1 and StGH2 genes were over-expressed in potato driven by the tuber specific promoter gbss, as described in example 5. A mutated AHAS gene was used as selection marker yielding tolerance to the Imazamox herbicides. Two potato varieties were transformed, Desiree and AM99-2003 a transgenic high amylose line with a 40% decrease in starch content compared to its mother variety. The transformed lines over-expressing StGH1 and StGH2 were selected as described in example 6. The gene expression levels were analysed with real-time PCR, see example 7 and table 3. The over-expression of the genes StGH1 and StGH2 resulted in an 2 two10 times increase in gene compared to their mother variety. Furthermore the lines over-expressing StGH1 and StGH2 showed an increase in dry matter of up to 36 % as described in example 8 and table 5.

The over-expression of StGH1 and StGH2 in transgenic potato plants producing amylose type starch resulted in an increased dry matter content, which means an increased amylose content as no amylopectine is produced, see example 8, 9 and 11. The amylose biosynthesis enhancing protein according to the invention comprises the amino acid sequence SEQ ID NO 2 or 4 or a protein which comprises a sequence derived from SEQ ID NO 2 or 4, which is at least 50%, preferably at least 60%, more preferably at least 70%, more preferably at least 80%, still more preferably at least 90%, most preferably at least 95%, identical at the amino acid level to the sequence SEQ ID NO 2 or 4 and has the property of an amylose biosynthesis enhancing protein. This amylose biosynthesis enhancing protein may also be prepared by artificial variations starting from the SEQ ID NO 2 or 4, for example by substitution, insertion or deletion of amino acids.

10

30

40

5

Such a protein could also be used to increase the production of starch or amylopectin in non-transgenic or transgenic plants.

The term "substitution" in the specification means the replacement of one or more amino acids by one or more amino acids. Preference is given to carrying out "conservative" replacements in which the amino acids replaced has a property similar to that of the original amino acid, for example replacement of Glu by Asp, Gln by Asn, Val by Ile, Leu by Ile, Ser by Thr.

"Deletion" is the replacement of an amino acid or amino acids by a direct bond. Preferred positions for deletions are the polypeptide termini and the junctions between the individual protein domains.

"Insertions" are insertions of amino acids into the polypeptide chain, with a direct bond formally being replaced by one or more amino acids.

"Identity" between two proteins means the identity of the amino acids over the in each case entire length of the protein, in particular the identity which is calculated by comparison with the aid of the Vector NTI Suite 7.1 Software of the company informax (USA) using the Clustal W method (Thompson, JD et al., Nucleic Acid Research, 22 (22):4673-4680, 1994)

with the parameters set as follows:

35 Multiple alignment parameter:

Gap opening penalty 15
Gap extension penalty 6.66
Gap separation penalty range 8
Gap separation penalty on

BASF Plant Science GmbH

20020884 84:81 x18M.7 21iszsanstam3

% identity for alignment delay 40
Residue specific gaps on
Hydrophilic residue gap off
Transition weighing 0

5

Pairwise alignment parameter:

10	FAST algorithm K-tuple size	off 2
	Gap penalty	5
	Window size	4
	Number of best diagonals	4

Accordingly, a protein which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4 means a protein which, when comparing its sequence with the sequence SEQ ID NO 2 or 4, is at least 50% identical, in particular according to the above program algorithm using the above set of parameters.

Further natural examples of genes coding for an amylose enhancing protein according to the invention can readily be found, for example, in various organisms, in particular in plants, whose genomic sequence is known by comparing the identity of the amino acid sequences or of the corresponding back-translated nucleic acid sequences from data-bases with the sequence of SEQ ID NO 2 or 4, in particular according to the above program algorithm using the above set of parameters.

In the completed genome sequence of Arabidopsis thaliana, five putitative coding sequences can be deduced by searching for exon/intron boundaries and comparing with back translated sequences of SEQ ID NO 2 or 4.

30

The following nucleic acid sequences of Arabidopsis thallana SEQ ID NO 5, SEQ ID NO 7, SEQ ID NO 9, SEQ ID NO 11 and SEQ ID NO 13 could be used to carry out the invention and are coding for the amylose biosynthesis enhancing proteins SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12 and SEQ ID NO 14.

35

Furthermore the following nucleic acid sequences or ESTs can be used in order to identify and clone genes coding for an amylose biosynthesis enhancing protein from plant organisms:

Tomato ESTs from GenBank: AW216407, BE450055, BF097262, BE450557, BF097173

Wheat ESTs from GenBank: BJ292476, BJ278875, BJ283925, BE442966, CA666180, BQ483228

Maize EST from GenBank: BG319971

Rice ESTs from GenBank: AL606633, CA752890, BI813265

10

15

Natural examples of amylose biosynthesis enhancing proteins and the corresponding genes can furthermore readily be found in various organisms, in particular plants, whose genomic sequence is unknown by hybridization techniques in a manner known per se, for example starting from the nucleic acid sequences SEQ ID NO 1 or SEQ ID NO 3 or any of the SEQ ID NO 5, 7, 9, 11 or 13 or any of the EST sequences described above.

The hybridization may be carried out under moderate (low stringency) or, preferably, under stringent (high stringency) conditions.

20

Such hybridization conditions are described, inter alia, in Sambrook, J., Fritsch, E.F., Maniatis, T., in: Molecular Cloning (A Laboratory Manual), 2nd edition, Cold Spring Harbor Laboratory Press, 1989, pages 9.31-9.57 or in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6.

25

By way of example, the conditions during the washing step may be selected from the range of conditions which is limited by those with low stringency (with 2X SSC at 50°C) and those with high stringency (with 0.2X SSC at 50°C, preferably at 65°C) (20X SSC: 0.3 M sodium citrate, 3 M sodium chloride, pH 7.0).

30

· 35

In addition, the temperature may be raised during the washing step from moderate conditions at room temperature, 22°C, to stringent conditions at 65°C.

Both parameters, salt concentration and temperature, may be varied simultaneously and it is also possible to keep one of the two parameters constant and to vary only the other one. It is also possible to use denaturing agents such as, for example, formamide or SDS during hybridization. In the presence of 50% formamide, the hybridization is preferably carried out at 42°C.

40 Some exemplary conditions for hybridization and washing step are listed below:

- (1) hybridization conditions with, for example
- (i) 4X SSC at 65°C, or

- (ii) 6X SSC at 45°C, or
- (iii) 6X SSC at 68°C, 100 mg/ml denatured fish sperm DNA, or
- (iv) 6X SSC, 0.5% SDS, 100 mg/ml denatured fragmented salmon sperm 10 DNA at 68°C, or
 - (v) 6X SSC, 0.5% SDS, 100 mg/ml denatured fragmented salmon sperm DNA, 50% formamide at 42°C, or

15

- (vi) 50% formamide, 4X SSC at 42°C, or
- (vii) 50% (vol/vol) formamide, 0.1% bovine serum albumin, 0.1% Ficoll, 0.1% polyvinylpyrrolidone, 50 mM sodium phosphate buffer pH
- 6.5, 750 mM NaCl, 75 mM sodium citrate at 42°C, or 20
 - (viii) 2X or 4X SSC at 50°C (moderate conditions), or
 - (ix) 30 to 40% formamide, 2X or 4X SSC at 42°C (moderate conditions).

25

- (2) Washing steps of 10 minutes each with, for example
- (i) 0.015 M NaCl/0.0015 M sodium citrate/0.1% SDS at 50°C, or
- 30 (ii) 0.1X SSC at 65°C, or
 - (iii) 0.1X SSC, 0.5% SDS at 68°C, or
 - (iv) 0.1X SSC, 0.5% SDS, 50% formamide at 42°C, or

- (v) 0.2X SSC, 0.1% SDS at 42°C, or
- (vi) 2X SSC at 65°C (moderate conditions).

Preferred proteins with amylose biosynthesis enhancing activity are proteins from plants, cyanobacteria, mosses or algae, particular preferred from plants. A particular preferred protein comprises the amino acid sequence SEQ ID NO 2 or 4.

- If, for example, the protein is to be expressed in a plant, it is frequently advantageous to use the codon usage of said plant for backtranslation and resynthesis of the gene according to codon usage of said plant.
- The invention further relates to nucleic acids encoding an amylose biosynthesis enhancing protein according to the invention. All of the nucleic acids mentioned in the specification may be, for example, a RNA sequence, DNA sequence or cDNA sequence.
- Suitable nucleic acid sequences can be obtained, for example, by back-translating the polypeptide sequence according to the genetic code. For this, preference is given to using those codons which are used frequently according to the organism-specific codon usage. The codon usage can be readily determined on the basis of computer analyses of other known genes of the organisms in question.
- All of the above-mentioned genes coding for an amylose biosynthesis enhancing protein can furthermore be prepared in a manner known per se from the nucleotide building blocks by chemical synthesis, for example by fragment condensation of individual overlapping complementary nucleic acid building blocks of the double helix. The chemical synthesis of oligonucleotides may be carried out, for example, in a known manner according to the phosphoramidite method (Voet, Voet, 2nd edition, Wiley Press New York, pages 896-897). The annealing of synthetic oligonucleotides and filling-in of gaps with the aid of the Klenow fragment of DNA polymerase and ligation reactions and also general cloning methods are described in Sambrook et al. (1989), Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press.

Genes coding for this function may be integrated in the plant chromosomes and upon expression utilize a transit peptide to localise to plastids which is the organelle where amylose biosynthesis takes place or be integrated directly into the plastid genome and thereby surpass the need for the localisation signal. The genes may be expressed constitutively or organ specific. For organ specific expression, promoters with tuber specific expression is preferable in potatoes while in cereals as maize or wheat a endosperm specific expression would be preferred to achieve a high degree of expression in organs where storage starch is accumulated. When transformed to the plastid genome then specific regulatory elements suitable for that organelle apply.

30

15

20

25

30

35

40-

20020884

The genes of this invention may be used in combination with other genes that can be situated on the same gene construct or transferred and combined by co-transformation or super transformation. Genes and traits that are of interest to combine with the genes of the instant invention are agronomic or input trait such as herbicide tolerance, disease and pest resistance or stress tolerance but could also be output traits such as starch structure modification or yield. Genes and traits used in combination with the genes described in the invention could be for adding a function that is not present in the modified plant species or over-expressing a function that is already present or inhibiting a function by the use of antisense, RNAi or antibodies.

The invention may be used to increase the amylose content in potato tubers but would in its context not be limited to potatoes but would be applicable to other starch producing and storing plants such as e.g. corn, cassava, wheat, barley, oat and rice.

The described invention is particularly suited for eliminating a lower starch content associated with increased amylose content in different plants where the number of α-1,4-glucan chain non-reducing ends is greatly reduced due to the reduction or elimination of α-1,6 branch formation. Amylopectin is an extremely efficient structure, as is glycogen, for polysaccharide production since it is very branched and thus contains as many points accessible for starch synthesis as there are non-reducing ends. Starch that is mainly composed of amylose, contains much fewer branches and thus the biosynthetic capacity is reduced. In order to enhance starch biosynthesis when there is no amylopectin production, expression of genes as described in the present invention, could for example form new primers that can replace amylopectin as a source for starch biosynthesis capacity and thereby reduce or eliminate the lost capacity for starch synthesis. To further illustrate the situation the degree of branching in ordinary potato starch is approximately 3.1% while in high amylose starch it is 0.3-1.0% depending on amylose content this decrease of branching and starch content is further associated with an increase in glucose and fructose content.

The increased amylose content and thereby solids content is also advantageous for the processing properties in various applications such as for french fries, potato crisps and other potato based products. In addition to an increased solid content, the inserted genes SEQ ID NO 1 or 3 of the present invention result in the transformation of excess sugars into α -1,4-glucan chains and thereby reducing browning of fried potato products. Maillard reaction, in which amino acids react with free sugars.

Furthermore

(i) any gene of plant origin with the described activity can be used for increasing amylose content and solids

25

35

- (ii) the genes can be controlled by any regulating promoter element functional in plant.
- (iii) any starch producing crop of any variety can be transformed with the described genes.
- 5 (iiii) any plant transformation method can be used.
 - (iiiii) any binary vector can be used for the insertion of the described genes.
 - (iiiiii) the described genes can be combined with any other desired transgenically inserted traits.
- The invention further relates to a method for producing amylose by culturing plants which have, compared to a wild type or a genetically modified plant producing already amylose type starch, an increased amylose biosynthesis activity, said proteins comprising the amino acid sequence SEQ ID NO 2 or 4 or a sequence which is derived from one of these sequences by substitution, insertion or deletion of amino acids and which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4.

Increased amylose biosynthesis activity compared to the wild type or transgenic line means that the amount of amylose formed is increased by the amylose biosynthesis enhancing protein in comparison with the wild type or transgenic line.

This increase in amylose biosynthesis activity is preferably at least 5%, further preferably at least 10%, further preferably at least 20%, further preferably at least 50%, more preferably at least 100%, still more preferably at least 200%, in particular at least 500%, of the protein activity of the wild type or transgenic line.

A "wild type" means the corresponding genetically unmodified starting plant. This plant is preferably Solanum tuberosum.

Depending on the context, the term "plant" means a wild type starting plant or a genetically modified starting plant.

"Transgenic plant" or "genetically modified plant" means that the plant contains an additional inserted gene segment that may be foreign or endogenous to the plant species, additional genes or additional gene fragments in sense and/or antisense orientation to a suitable promoter corresponding to the following polypeptides and showing enzymatic activity of a starch branching enzyme I, a starch branching enzyme II and/or the amylose biosynthesis enhancing protein as specified in SEQ ID NO 1 or 3 or polynucleotides having at least 60 % sequence identity thereof.

20

25

30

35

15

"Amylose type starch" means that the amylose content of the starch is increased compared to the amylose content of starch produced by wild type plants especially wild type potato plants.

The amylose biosynthesis activity may be increased in various ways, for example by eliminating inhibiting regulatory mechanisms at the translation and protein levels or by increasing the gene expression of a nucleic acid encoding a amylose biosynthesis enhancing protein compared to the wild type or transgenic plant, for example by inducing a gene encoding the amylose biosynthesis enhancing protein via activators or by introducing into the plant nucleic acids encoding a amylose biosynthesis enhancing protein.

According to the invention, increasing the gene expression of a nucleic acid encoding a amylose biosynthesis enhancing protein could also mean manipulating the expression of the endogenous amylose biosynthesis enhancing protein intrinsic to the plant, in particular in potato plants. This may be achieved, for example, by modifying the promoter DNA sequence of genes encoding an amylose biosynthesis enhancing protein. Such a modification which leads to a modified or preferably increased rate of expression of at least one endogenous gene encoding an amylose biosynthesis enhancing protein may be carried out by deleting or inserting DNA sequences.

It is also possible to modify expression of one or more endogenous amylose biosynthesis enhancing protein by applying exogenous stimuli. This may be carried out by particular physiological conditions, i.e. by applying foreign substances.

Furthermore, it is possible to achieve a modified or increased expression of at least one endogenous gene encoding an amylose biosynthesis enhancing protein by the interaction of a regulatory protein which is modified or is not present in the untransformed plant.

In a preferred embodiment, the amylose biosynthesis enhancing protein activity is increased compared to the wild type or transgenic plant by increasing the gene expression of a nucleic acid encoding an amylose biosynthesis enhancing protein, said amylose biosynthesis enhancing protein comprising the amino acid sequence SEQ ID NO 2 or 4 or a sequence which is derived from said sequences by substitution, insertion or deletion of amino acids and which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4.

In the case of genomic nucleic acid sequences encoding an amylose biosynthesis enhancing protein from eukaryotic sources, which contain introns, preferably already

20

25.

35

processed nucleic acid sequences such as the corresponding cDNAs are to be used, if the host organism is unable to or cannot be enabled to express the corresponding amylose biosynthesis enhancing protein.

In this preferred embodiment, the transgenic plant of the invention thus contains, compared to the wild type or transgenic plant, at least one further gene encoding an armylose biosynthesis enhancing protein. In this preferred embodiment, the genetically modified plant of the invention has accordingly at least one transgenic endogenous or exogenous nucleic acid encoding an amylose biosynthesis enhancing.

Suitable and preferred nucleic acids are described above. In a particularly preferred embodiment, a nucleic acid comprising the sequence SEQ ID NO 1 or 3 is introduced into the plant.

According to the invention, organisms means preferably eukaryotic organisms, such as, for example, yeasts, algae, mosses, fungi or plants, which are capable of producing starch or amylose, either as wild type or enabled by genetic modification. Preferred organisms are photosynthetically active organisms such as, for example, plants which, even as a wild type, are capable of producing starch or amylose type starch.

Particularly preferred organisms are potato plants.

The present invention furthermore relates to the use of proteins comprising the amino acid sequence SEQ ID NO 2 or 4 or a sequence which is derived from this sequence by substitution, insertion or deletion of amino acids and which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4 and having amylose biosynthesis enhancing activity.

The present invention further relates to the use of nucleic acids SEQ ID NO 1 or 3 or one of the SEQ ID NOs 5, 7, 9, 11 or 13 encoding proteins having an amylose biosynthesis enhancing activity in plants.

The transgenic organisms, in particular plants, are preferably prepared by transforming the starting organisms, in particular plants, with a nucleic acid construct containing the above-described nucleic acid, encoding an amylose biosynthesis enhancing protein which is functionally linked to one or more regulatory signals ensuring transcription and translation in said organisms.

These nucleic acid constructs in which the coding nucleic acid sequence is functionally linked to one or more regulatory signals ensuring transcription and translation in organisms, in particular in plants, are also referred to as expression cassettes herein below.

Accordingly, the invention further relates to nucleic acid constructs, in particular to 5 nucleic acid constructs functioning as expression cassette, which comprise a nucleic acid encoding an amylose biosynthesis enhancing protein which is functionally linked to one or more regulatory signals ensuring transcription and translation in organisms, in 10

The regulatory signals preferably comprise one or more promoters ensuring transcription and translation in organisms, in particular in plants.

The expression cassettes include regulatory signals, i.e. regulatory nucleic acid sequences, which control expression of the coding sequence in the host cell. According 15 to a preferred embodiment, an expression cassette comprises upstream, i.e. at the 5' end of the coding sequence, a promoter and downstream, i.e. at the 3' end, a polyadenylation signal and, where appropriate, further regulatory elements which are operatively linked to the coding sequence for at least one of the above-described genes located in between. Operative linkage means the sequential arrangement of promoter, 20 coding sequence, terminator and, where appropriate, further regulatory elements in such a way that each of the regulatory elements can properly carry out its function in the expression of the coding sequence.

25 When the organism used is a plant, the nucleic acid constructs and expression cassettes of the invention preferably contain a nucleic acid encoding a plastid transit peptide ensuring localisation in plastids.

The preferred nucleic acid constructs, expression cassettes and vectors for plants and methods for preparing transgenic plants and also the transgenic plants themselves are 30 described in examples 2 to 6 below.

The sequences preferred for operative linkage, but not limited thereto, are targeting sequences for ensuring subcellular localisation to plastids such as amyloplasts or chloroplasts but could also mean in the apoplasts, in the vacuole, in the mitochondrion, in the endoplasmic reticulum (ER), in the nucleus, in elaioplasts or in other compartments and translation enhancers such as the tobacco mosaic virus 5'-leader sequence (Gallie et al., Nucl. Acids Res. 15 (1987), 8693-8711).

35

40

20020884

A suitable promoter of the expression cassette is in principle any promoter which is able to control the expression of foreign genes in plants.

"Constitutive" promoter means those promoters which ensure expression in numerous, preferably all, tissues over a relatively long period of plant development, preferably during the entire plant development.

Preference is given to using, in particular, a promoter from plants or a promoter originating from a plant virus. Preference is in particular given to the promoter of the 35S transcript of the CaMV cauliflower mosaic virus (Franck et al. (1980) Cell 21:285-294; Odell et al. (1985) Nature 313:810-812; Shewmaker et al. (1985) Virology 140:281-288; Gardner et al. (1986) Plant Mol Biol 6:221-228) or the 19S CaMV promoter (US 5,352,605; WO 84/02913; Benfey et al. (1989) EMBO J 8:2195-2202).

Another suitable constitutive promoter is the Rubisco small subunit (SSU) promoter (US 4,962,028), the leguminB promoter (GenBank Acc. No. X03677), the Agrobacterium nopaline synthase promoter, the TR double promoter, the agrobacterium OCS (octopine synthase) promoter, the ubiquitin promoter (Holtorf S et al. (1995) Plant Mol Biol 29:637-649), the ubiquitin 1 promoter (Christensen et al. (1992) Plant Mol Biol 18:675-689; Bruce et al. (1989) Proc Natl Acad Sci USA 86:9692-9696), the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (US 5,683,439), the promoters of the vacuolar ATPase subunits or the promoter of a proline-rich wheat protein (WO 91/13991), the Pnit promoter (Y07648.L, Hillebrand et al. (1998), Plant. Mol. Biol. 36, 89-99, Hillebrand et al. (1996), Gene, 170, 197-200) and other promoters of genes whose constitutive expression in plants is known to the skilled worker.

The expression cassettes may also contain a chemically inducible promoter (review: Gatz et al. (1997) Annu Rev Plant Physiol Plant Mol Biol 48:89-108) which may be used to control expression of the amylose biosynthesis enhancing protein gene in the plants at a particular time. Promoters of this kind, such as, for example, the PRP1 promoter (Ward et al. (1993) Plant Mol Biol 22:361-366), salicylic acid-inducible promoter (WO 95/19443), a benzenesulfonamide-inducible promoter (EP 0 388 186), a tetracycline-inducible promoter (Gatz et al. (1992) Plant J 2:397-404), an abscisic acid-inducible promoter (EP 0 335 528) and an ethanol- or cyclohexanone-inducible promoter (WO 93/21334), may likewise be used.

Further examples of suitable promoters are fruit ripening-specific promoters such as, for example, the fruit ripening-specific promoter from tomato (WO 94/21794, EP 409 625). Development-dependent promoters partly include the tissue-specific promoters, since individual tissues are naturally formed in a development-dependent manner.

20020884

Furthermore, preference is given in particular to those promoters which ensure expression in tissues or parts of the plant, in which, for example, biosynthesis of starch or amylose or of the precursors thereof takes place. Preference is given, for example, to promoters with specificities for leaves, stems, roots, seeds and tubers.

Seed-specific promoters are, for example, the phaseoline promoter (US 5,504,200; Bustos MM et al. (1989) Plant Cell 1(9):839-53), the promoter of the 2S albumin gene (Joseffson LG et al. (1987) J Biol Chem 262:12196-12201), the legumin promoter (Shirsat A et al. (1989) Mol Gen Genet 215(2): 326-331), the USP (unknown seed protein) promoter (Bäumlein H et al. (1991) Mol Gen Genet 225(3):459-67), the promoter of the napin gene (US 5,608,152; Stalberg K et al. (1996) L Planta 199:515-519), the sucrose-binding protein promoter (WO 00/26388) and the legumin B4 promoter (LeB4; Bäumlein H et al. (1991) Mol Gen Genet 225: 121-128; Baeumlein et al. (1992) Plant Journal 2(2):233-9; Fiedler U et al. (1995) Biotechnology (NY) 13(10):1090f), the Arabidopsis oleosin promoter (WO 98/45461), the Brassica Bce4 promoter (WO 91/13980) and the vicillin promoter (Weschke et al. 1988, Biochem. Physiol. Pflanzen 183, 233-242; Bāumlein H et al. (1991) Mol Gen Genet 225(3):459-67).

- Further suitable seed-specific promoters are those of the genes coding for high molecular weight glutenine (HMWG), gliadin, branching enzyme, ADP glucose pyrophosphatase (AGPase) and starch synthase. Preference is further given to promoters which allow seed-specific expression in monocotyledons such as e.g. corn, barley, wheat, rye, rice, etc. It is also possible to use advantageously the promoter of the lpt2 or lpt1 gene (WO 95/15389, WO 95/23230) or the promotors described in WO 99/16890 (promoters of the hordein gene, the glutelin gene, the oryzin gene, the prolamin gene, the gliadin gene, the glutelin gene, the kasirin gene and the secalin gene).
- Examples of tuber-, storage root- or root-specific promoters are the patatin promoter class I (B33), the potato cathepsin D inhibitor promoter and the potato granular bound starch synthase (GBSS) promoter as described in EP-A 0 921 191.

Examples of leaf-specific promoters are the cytosolic FBPase promoter from potato (WO 97/05900), the rubisco (ribulose-1,5-bisphosphate carboxylate) SSU (small sub-unit) promoter and the potato ST-LSI promoter (Stockhaus et al. (1989) EMBO J 8:2445-2451).

Further promoters suitable for expression in plants have been described (Rogers et al. (1987) Meth in Enzymol 153:253-277; Schardl et al. (1987) Gene 61:1-11; Berger et al. (1989) Proc Natl Acad Sci USA 86:8402-8406).

- The site of starch and amylose biosynthesis in potato plants is the amyloplast. Therefore amyloplast-specific targeting and activity of the gene products of the inventivenucleic acids SEQ ID NO 1 or 3 encoding an amylose biosynthesis enhancing protein is desirable.
- The expression may also take place in a tissue-specific manner in all parts of the plant. 10

A further preferred embodiment therefore relates to a tuber-specific expression of the nucleic acids SEQ ID NO 1 or 3.

- In addition, a constitutive expression of the gene encoding an amylose biosynthesis 15 enhancing protein is advantageous. On the other hand, however, an inducible expression of this gene may also be desirable.
- An expression cassette is preferably prepared by fusing a suitable promoter to an above-described nucleic acid encoding an amylose biosynthesis enhancing protein 20 and, preferably, to a nucleic acid which has been inserted between promoter and nucleic acid sequence and which codes for an amyloplast-specific transit peptide and also to a polyadenylation signal according to familiar recombination and cloning techniques as described, for example, in T. Maniatis, E.F. Fritsch and J. Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 25 (1989) and in T.J. Silhavy, M.L. Berman and L.W. Enquist, Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1984) and in Ausubel, F.M. et al., Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley-Interscience (1987). 30

Particular preference is given to inserted nucleic acid sequences which ensure targeting in the amyloplasts.

It is also possible to use an expression cassette in which the nucleic acid sequence encodes an amylose biosynthesis enhancing protein fusion protein, one part of the 35 fusion protein being a transit peptide which controls translocation of the polypeptide. Preference is given to amyloplast-specific transit peptides which, after translocation of amylose biosynthesis enhancing protein into the amyloplasts, are enzymatically cleaved off the amylose biosynthesis enhancing protein part.

10

15

20

25

30

35

Particular preference is given to the transit peptide which is derived from the Nicotiana tabacum plastid transketolase or from another transit peptide (e.g. the transit peptide of the rubisco small subunit or of ferredoxin NADP oxidoreductase and also of isopentenyl pyrophosphate isomerase-2) or from its functional equivalent.

Further examples of a plastid transit peptide are the transit peptide of the plastid isopentenyl pyrophosphate isomerase-2 (IPP-2) from Arabidopsis thaliana and the transit peptide of the ribulose bisphosphate carboxylase small subunit (rbcS) from pea (Guerineau, F, Woolston, S, Brooks, L, Mullineaux, P (1988) An expression cassettte for targeting foreign proteins into the chloroplasts. Nucl. Acids Res. 16: 11380).

Plant genes of the invention which encode a plant amylose biosynthesis enhancing protein may already contain the nucleic acid sequence which encodes a plastid transit peptide. In this case, a further transit peptide is not required. For example, the Solanum tuberosum sequences of the amylose biosynthesis enhancing protein of the invention SEQ ID NO 1 or 3 contain already a transit peptide sequence.

The nucleic acids of the invention may be prepared synthetically or obtained naturally or comprise a mixture of synthetic and natural nucleic acid components and may also be composed of various heterologous gene sections of various organisms.

As described above, preference is given to synthetic nucleotide sequences with codons which are preferred by plants. These codons which are preferred by plants may be determined from codons which have the highest frequency in proteins and which are expressed in most of the interesting plant species.

When preparing an expression cassette, it is possible to manipulate various DNA fragments in order to obtain a nucleotide sequence which expediently can be read in the correct direction and is provided with a correct reading frame. The DNA fragments may be linked to one another by attaching adaptors or linkers to said fragments.

It is furthermore possible to use manipulations which provide appropriate restriction cleavage sites or which remove excess DNA or restriction cleavage sites. In those cases for which insertions, deletions or substitutions such as, for example, transitions and transversions are suitable, in vitro mutagenesis, primer repair, restriction or ligation can be used.

Preferred polyadenylation signals are polyadenylation signals functional in plants, exemplified by those which correspond essentially to T-DNA polyadenylation signals from Agrobacterium tumefaciens, in particular of the T-DNA gene 3 (octopine synthase)

25

or OCS terminator, the complete sequence of the Ti plasmid pTiACH5 (Gielen et al., EMBO J. 3, 835 –846(1984) or functional equivalents.

The invention further relates to the use of the nucleic acids SEQ ID NO 1 or 3 for increasing the starch or amylose content in plants, e.g. potato plants which, as wild type, are capable of producing starch or amylose, see examples 2-12.

The invention is not limited to the over-expression of the nucleic acid sequences SEQ ID NO 1 or SEQ ID NO 3 in plants especially potato plants.

The over-expression of both nucleic acid sequences SEQ ID NO 1 and 3 can be used for enhancing amylose biosynthesis. Constructs containing the nucleic acids SEQ ID NO 1 and SEQ ID NO 3 can also be used for increasing the starch content or the amylopectin content in plants. These constructs can be made on the same T-DNA driven by one promoter each. These constructs can also be made on the same T-DNA in tandem driven by the same promoter. These constructs can also be transformed using more than one construct, either at the same time (co-transformation) or in different transformation events.

20 The above-described proteins and nucleic acids may be used for producing starch or amylose in transgenic plants.

The transfer of foreign genes into the genome of an organism, in particular of a plant, is referred to as transformation.

For this purpose, methods known per se for transforming plants and regenerating plants from plant tissues or plant cells can be used, in particular in plants, for transient or stable transformation, e.g. as described in example 2.

Suitable methods for the transformation of plants are the protoplast transformation by polyethylene glycol-induced DNA uptake, the biolistic method using the gene gun - also known as particle bombardment method, electroporation, the incubation of dry embryos in a DNA-containing solution, microinjection and the above-described Agrobacterium-mediated gene transfer. Said methods are described, for example, in B. Jenes et al.,
 Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, edited by S.D. Kung and R. Wu, Academic Press (1993), 128-143 and in Potrykus, Annu. Rev. Plant Physiol. Plant Molec. Biol. 42 (1991), 205-225).

The construct to be expressed is preferably cloned into a vector which is suitable for transforming Agrobacterium tumefaciens, for example pBin19 (Bevan et al., Nucl. Acids Res. 12 (1984), 8711) or preferably pSUN2 (WO 02/00900).

Accordingly, the invention furthermore relates to vectors containing the abovedescribed nucleic acids, nucleic acid constructs or expression cassettes.

Agrobacteria which have been transformed with an expression cassette can be used in a known manner for the transformation of plants, for example by bathing injured leaves or leaf sections in an agrobacteria solution and then culturing them in suitable media.

Apart from in plants, the expression cassette may also be used for transforming bacteria, in particular cyanobacteria, mosses, yeasts, filamentous fungi and algae.

- Genetically modified plants, also referred to as transgenic plants herein below, are preferably prepared by cloning the fused expression cassette which expresses a amylose biosynthesis enhancing protein into a vector, for example pBin19, which is suitable for transforming Agrobacterium tumefaciens.
- Agrobacteria which have been transformed with such a vector may then be used in a known manner for the transformation of plants, in particular of crop plants, for example by bathing injured leaves or leaf sections in an agrobacteria solution and then culturing them in suitable media.
- The transformation of plants by agrobacteria is described, inter alia, in F.F. White, Vectors for Gene Transfer in Higher Plants; in Transgenic Plants, Vol. 1, Engineering and Utilization, edited by S.D. Kung and R. Wu, Academic Press, 1993, pp. 15-38.
 Transgenic plants which contain a gene for expression of a nucleic acid encoding a amylose biosynthesis enhancing protein, which has been integrated into the expression cassette, can be regenerated in a known manner from the transformed cells of the injured leaves or leaf sections.

A host plant is transformed with a nucleic acid SEQ ID NO 1 or 3 encoding an amylose biosynthesis enhancing protein by incorporating an expression cassette as insertion into a recombinant vector whose vector DNA comprises additional functional regulatory signals, for example sequences for replication or integration. Suitable vectors are described inter alia, in Methods in Plant Molecular Biology and Biotechnology (CRC Press), chapter 6/7, pp. 71-119 (1993).

By way of example, the plant expression cassette may be incorporated into a derivative of the transformation vector pBin-19 with 35s promoter (Bevan, M., Nucleic Acids Research 12: 8711-8721 (1984).

Using the above-cited recombination and cloning techniques, it is possible to clone the expression cassettes into suitable vectors for maintenance and propagation of genetic material for example in E. coli. Suitable cloning vectors are, inter alia, pBR322, pUC series, M13mp series and pACYC184. Particularly suitable are binary vectors which can replicate both in E. coli and in agrobacteria.

10

The invention therefore further relates to the use of the above-described nucleic acids or of the above-described nucleic acid constructs, in particular of the expression cassettes, for preparing genetically modified plants or for transforming plants, plant cells, plant tissues or parts of plants.

15

25

30

35

40

The use is preferably aimed at increasing the starch or amylose content of the plant, of the tubers or in other parts of the plant.

The use is most preferably aimed at increasing the starch or amylose content of wildtype or transgenic potato plants and especially the tubers of wild-type or transgenic potato plants.

Accordingly, the invention further relates to a method for preparing genetically modified plants by introducing an above-described nucleic acid or an above-described nucleic acid construct into the genome of the starting organism.

The invention further relates to the genetically modified organisms, the genetic modification increasing the activity of an amylose biosynthesis enhancing protein compared to a wild type or transgenic plant and the amylose biosynthesis enhancing protein comprising the amino acid sequence SEQ ID NO 2 or 4 or a sequence which is derived from this sequence by substitution, insertion or deletion of amino acids and which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4.

As illustrated above, the amylose biosynthesis enhancing protein activity is increased compared to the wild type or transgenic plant preferably by increasing the gene expression of a nucleic acid encoding an amylose biosynthesis enhancing protein.

In a further preferred embodiment, gene expression of a nucleic acid encoding a amylose biosynthesis enhancing protein is increased, as illustrated above, by introducing nucleic acids encoding an amylose biosynthesis enhancing protein into the organism and thus by over-expressing nucleic acids encoding an amylose biosynthesis enhancing protein.

Such transgenic plants, their propagation material and their plant cells, plant tissues, plant parts or tubers are a further subject of the present invention.

Genetically modified plants of the invention, which have an increased starch or amylose content and which can be consumed by humans and animals, can also be used as
food- or feedstuffs or as feed and food supplements, for example directly or after processing known per se. The genetically modified plants may furthermore be used for
producing starch or amylose-containing extracts of said plant and/or for producing feed
and food supplements.

The invention further relates to:

15

10

- A polynucleotide that encodes a polypeptide of SEQ ID NO 1 or 3.
- II. A polynucleotide comprising at least 30 contiguous bases of SEQ ID NO 1 or 3.
- 20 III. A polynucleotide having at least 60 % sequence identity to SEQ ID NO 1 or 3, wherein the identity is based on the entire coding sequence.
 - IV. A polynucleotide having at least 60 % sequence identity to SEQ ID NO 1 or 3, wherein the % sequence identity is based on the entire sequence.

25

V. A polynucleotide which selectively hybridizes, under stringent conditions and a wash in 2 X SSC at 50 °C, to a hybridization probe derivable from the polynucleotide sequence as set forth in SEQ ID NO 1 or 3, or from the genomic sequence.

30

- VI. A polynucleotide complementary to a polynucleotide of V.
- VII. The polynucleotide of I, wherein the starch or amylose biosynthesis enhancing polynucleotide is from Solanum tuberosum.

- VIII. The polynucleotide of I encoding a polypeptide, which after over-expression in a plant cell increases the starch or amylose content.
- IX. The polynucleotide of I in antisense orientation, which after expression in a plant cell decreases the starch or amylose content.

- X. A vector comprising at least one polynucleotide of I.
- XI. An expression cassette comprising at least one polynucleotide of I operably linked to a promoter, wherein the polynucleotide is in sense or antisense orientation.
 - XII. A host cell which is introduced with at least one expression cassette of X.
- 10 XIII. The host cell of XI that is a plant cell.
 - XIV. A transgenic plant comprising at least one expression cassette of XI.
 - XV. The transgenic plant of XIII, wherein the plant is Solanum tuberosum.
- XVI. A tuber from the transgenic plant of XIV.

15

20

25

- XVII. An isolated protein comprising a member selected from the group consisting of:
 - a) a polypeptide comprising at least 10 contiguous amino acids of SEQ ID
 NO 2 or 4,
 - b) a polypeptide which is a plant amylose biosynthesis enhancing protein,
 - c) a polypeptide comprising at least 55 % sequence identity to SEQ ID NO 2 or 4, wherein the sequence identity is based on the entire sequence and has at least one epitope in common with an amylose biosynthesis enhancing protein.
 - a polypeptide encoded by a polynucleotide selected from SEQ ID NO 1 or 3,
 - e) a polypeptide of SEQ ID NO 2 or 4.
- 30 XVIII. The protein of XVII, wherein the polypeptide is catalytically active.
 - XIX. A ribonucleic acid sequence encoding the protein of XVIII.
- XX. A method for modulating the level of amylose biosynthesise enhancing protein in a plant, comprising:
 - a) stably transforming a plant cell with a polynucleotide coding for an amylose biosynthesis enhancing protein operably linked to a promoter, wherein the polynucleotide is in sense or antisense orientation
 - growing the plant cell under plant growing conditions to produce a regenerated plant capable of expressing the polynucleotide for a time sufficient

25

30

35

40

20020884

to modulate the level of amylose biosynthesis enhancing protein in the plant.

- XXI. The method of XX, wherein the polynucleotide coding for an amylose biosynthesis enhancing protein is selected from SEQ-ID NO 1 or 3.
 - XXII. The method of XX, wherein the plant is Solanum tuberosum.
- XXIII. The method of XX, wherein activity of the amylose biosynthesis enhancing protein is increased.
 - XXIV. A method for modulating the level of starch or amylose in a plant, comprising:
 - a) stably transforming a plant cell with a polynucleotide coding for an amylose biosynthesis enhancing protein operably linked to a promoter, wherein the polynucleotide is in sense or anti-sense orientation,
 - growing the plant cell under plant growing conditions to produce a regenerated plant capable of expressing the polynucleotide for a time sufficient to modulate level of starch or amylose in the plant.
- 20 XXV. A method for modulating the level of amylose in a plant, comprising:
 - a) stably transforming a plant cell with a polynucleotide encoding an amylose biosynthesis enhancing protein operably linked to a promoter, wherein the polynucleotide is in sense or anti-sense orientation.
 - growing the plant cell under plant growing conditions to produce a regenerated plant capable of expressing the polynucleotide for a time sufficient to modulate level of amylose in the plant.
 - XXVI. The method of XXIV wherein the polynucleotide coding for an amylose biosynthesis enhancing protein is selected from SEQ ID NO 1 or 3.

Some of the terms used further on in the specification are defined at this point.

"Enzymatic activity/activity assay": the term enzymatic activity describes the ability of an enzyme to convert a substrate into a product. In this context, both the natural substrate of the enzyme and a synthetic modified analog of the natural substrate can be used. The enzymatic activity can be determined in what is known as an activity assay via the increase in the product, the decrease in the starting material, the decrease or increase in a specific cofactor, or a combination of at least two of the aforementioned parameters as a function of a defined period of time.

10

15

25

30

35

40

"Functional equivalents" in the present context describe nucleic acid sequences which hybridize under standard conditions with the nucleic acid sequence encoding the amylose biosynthesis enhancing protein or portions of the nucleic acid sequence encoding the amylose biosynthesis enhancing protein, and which are capable of bringing about the expression of an enzymatically active plant amylose biosynthesis enhancing protein in a cell or an organism.

It is advantageous to use short oligonucleotides of a length between 10 to 50bp, preferably 15-40bp, for example of the conserved or other regions, which can be determined via comparisons with other related genes in a manner known to the skilled worker for the hybridization. Alternatively, it is also possible to use longer fragments of the nucleic acids according to the invention or the complete sequences for the hybridization. These standard conditions vary depending on the nucleic acid used, namely oligonucleotide, longer fragment or complete sequence, or depending on which type of nucleic acid, that is DNA or RNA, is being used for the hybridization. Thus, for example, the melting temperatures for DNA:DNA hybrids are approx. 10oC lower than those of DNA:RNA hybrids of equal length. Suitable hybridization conditions are described above.

A functional equivalent is furthermore also understood as meaning, in particular, natural or artificial mutations of the relevant nucleic acid sequences of the plant amylose biosynthesis enhancing protein and their homologs from other organisms which make possible the expression of the enzymatically active plant amylose biosynthesis enhancing protein in a cell or an organism.

Thus, the scope of the present invention also extends to, for example, those nucleotide sequences which are obtained by modification of the nucleic acid sequence of a amylose biosynthesis enhancing protein. The purpose of such a modification can be, for example, the insertion of further cleavage sites for restriction enzymes, the removal of excess DNA, or the addition of further sequences. Proteins which are encoded via said nucleic acid sequences should still maintain the desired functions, despite the deviating nucleic acid sequence.

The term functional equivalent may also refer to the protein encoded by the nucleic acid sequence in question. In this case, the term functional equivalent describes a protein whose amino acid sequence is up to a specific percentage identical with that of the amylose biosynthesis enhancing protein.

Functional equivalents thus encompass naturally occurring variants of the sequences described herein, and also artificial, for example chemically synthesized, nucleic acid

20020884

sequences adapted to the codon usage, or the amino acid sequences derived there-from.

In general, it can be said that functional equivalents independently of the amino acid sequence in question (encoded by a corresponding nucleic acid sequence) have in each case the enzymatic activity of a amylose biosynthesis enhancing protein.

"Reporter genes" encode readily quantifiable proteins. Using these genes, an assessment of transformation efficacy or of the site or time of expression can be made via growth, fluorescence, chemoluminescence, bioluminescence or resistance assay or via 10 photometric measurement (intrinsic color) or enzyme activity. Very especially preferred in this context are reporter proteins (Schenborn E, Groskreutz D. Mol. Biotechnol. 1999; 13(1):29-44) such as the "green fluorescence protein" (GFP) (Gerdes HH and Kaether C, FEBS Lett. 1996; 389(1):44-47; Chui WL et al., Curr. Biol. 1996, 6:325-330; Leffel SM et al., Biotechniques. 23(5):912-8, 1997), chloramphenicol acetyl transferase, 15 a luciferase (Giacomin, Plant Sci. 1996, 116:59-72; Scikantha, J. Bact. 1996, 178:121; Millar et al., Plant Mol. Biol. Rep. 1992 10:324-414), and luciferase genes, in general bgalactosidase or b-glucuronidase (Jefferson et al., EMBO J. 1987, 6, 3901-3907), the Ura3 gene, the liv2 gene, the 2-desoxyglucose-6-phosphate phosphatase gene, blactamase gene, the neomycin phosphotransferase gene, the hygromycin phos-20 photransferase gene, or the BASTA (= gluphosinate) resistance gene.

"Significant increase": referring to the enzymatic activity, is understood as meaning the increase in the enzymatic activity of the enzyme incubated with a candidate compound in comparison with the activity of an enzyme not incubated with the candidate compound, which lies outside an error in measurement.

"Substrate": Substrate is the compound which is recognized by the enzyme in its original function and which is converted into a product by means of a reaction catalyzed by the enzyme.

Preferably, the plant amylose biosynthesis enhancing protein is encoded by a nucleic acid sequence comprising

- 35 a) a nucleic acid sequence shown in SEQ ID NO 1 or 3; or
 - b) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO 2 or 4 by back translation; or

25

30 .

a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from a functional equivalent of the amino acid sequence shown in SEQ ID NO 2 or 4, which has an identity with SEQ ID NO 2 or 4 of at least 50%, by back translation.

5

10

20

25

The functional equivalent of SEQ ID NO 2 or 4 set forth in c) has an identity of at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57% preferably at least 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, and 70% more preferably 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85% most preferably at least 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID NO 2 or 4.

Example 1

15 Complementation study in yeast

Yeast contains two self-glycosylating proteins, Glg1p and Glg2p, which yield primers for the initiation of glycogen synthesis. For glycogen synthesis to take place in yeast it is required that either gene is functional. Yeast strain CC9, contain knock-out mutations for both genes and is therefore a null mutant regarding this specific biosynthetic function and is therefore unable to produce glycogen (Cheng, C. et al., Molecular and Cellular Biology (1995), 6632-6640). CC9 was used as a basis for complementation experiments with the isolated potato genes in order to validate their function by restoring glycogen biosynthesis in the CC9 strain. The potato genes were cloned in a yeast plasmid, pRS414 (Stratagene), and expressed with various yeast controlling elements such as Gal1, Adh1 and Glg2p promoters. CC9 was transformed by the resulting plasmids using LiCl and electroporation (Multiporator, Eppendorf). Transformed yeast colonies growing on appropriate media plates were screened by immersing in iodine solution. Wild type yeast producing glycogen is stained red brown by lodine while the null mutant CC9 is not stained. CC9 expressing the potato genes, StGH1 and StGH2, will stain red brown, when the isolated genes complement a glycogenin function in yeast and thus carry the desired function.

Example 2

35

30

Transformation method

Fully expanded leaves from in vitro propagated potato plants are diagonally cut in 2-4 pieces and precultivated on MC-plates for 2-3 days at 23-24°C.

Agrobacterium tumefaciens strain LBA4404 containing pHS1, pHS2, pHS3 or pHS4 are grown in YEB medium with 100µg rifampicin and 25µg/ml kanamycin over night on constant shaking (200 rpm) at 28°C.

The Agrobacterium culture is prepared for infection by dilution 1:20 with MS10 medium. The leaf explants are infected for 8-10 min in the bacterial solution and afterwards drained on filter paper for 5-20 seconds. The leaf segments are placed on the MS300 plates for 2 days co-cultivation under modest light at 23-24°C. At the end of co-cultivation the leaf segments are moved to M400 plates containing 400mg/l Claforan to suppress bacterial growth. After 4-5 days the explants are moved to selection medium MS400 supplemented with 400mg/l Claforan. For explants transformed with pHS1 and pHS2 50µM kanamycin was included in the media and for explants transformed with pHS3 and pHS4 0.5µM Imazamox was added to the media.

Leaf segments are transferred to fresh MS 400 selection medium every fortnight. The regenerated putative transgenic shoots are collected and cultivated on MS30 plates with 200mg/l Claforan aiming at shoot elongation.

When the shoots are 3-5 cm long, 1-2 cm are cut off and grown on microtuber medium in the dark at 25°C. After 2-5 weeks microtubers are produced. Putative transgenic plants are analysed for GUS expression in microtubers to determine the transformation efficiency.

MC plates	MS300
MS300 plates with 1.5-2 ml liquid MS100 medium and covered with one sterile filter paper	4.4g/l MS-medium 2mg/l naphtyl acetic acid 1mg/l 6-benzyl amino pyridine 3% (w/v) sucrose pH 5.2
MS10	MS400
4.4 g/l MS-medium (murashige and Skoog) 1% (w/v) sucrose pH 5. 8	4.4 g/l MS-medium 2mg/l zeatine 0.01mg/l naphtyl acetic acid mg/l gibberellic acid 10% (w/v) sucrose 400 mg/l claforan 0.5 µM Imazamox or 50 µM kanamycin pH5.8

PH 5,8	- N. G. W.
1	Mg/l gibberellic acid
	10% (w/v) sucrose
	400 mg/l claforan
·	0,5 μM Imazamox or 50 μM kanamacin
µM MS30	pH 5.8
MINI MIGSO	Microtuber medium
4.4 g/l MS-medium	
3% (w/v) sucrose	4.4 g/l MS- medium
pH 5.8	2.5 mg/l kinetin
p. 1 0.10	0.5 mg/l abscisic acid
	8% sucrose
MS100	200 mg/ claforan
4.4 mg/l MS-medium	
30g/i sucrose	
0.5 m/lg thiamin-HCl	
0.5 mg/l pyridoxin-HCl	
mg/l nicotinacid	
0.5 mg/l kinetin	1.
29.8 mg/l ferrous sulfate hepta hydrate	
mg/l 2,4-Dichlorophenoxyacetic acid	
g/i caseinhydrolysate	
H 5.2	<u>'</u>

Example 3

Transgenic plant AM 99-2003

5

High amylose potato lines can be produced for example by using antisense, RNAi or antibody technology that target the two starch branching enzymes starch branching enzyme 1 (SBE1) and starch branching enzyme 2 (SBE2).

- The high amylose potato line AM99-2003 is produced by inhibition of the starch branching enzyme activities in the mother variety Dinamo. Transformation is made with a construct of SBE1 and SBE2 in antisense orientation driven by the gbss promoter.
- pBluescript containing a 1620bp fragment of the 3'end of Sbe1 between EcoRV and Spel is cut open with Spel (blunt) and Xbal and ligated with a 1243bp Sstl (blunt) and

<

Xbal fragment of the 3'end of Sbe2. The Sbe2 and Sbe1 complex is cut out with EcoRV and Xbal and ligated to the Smal and Xbal opened up binary vector pHo3.1, see figure 8. The final vector is named pHAbe12A, see figure 9 and nucleic acid sequence SEQ ID NO 15. pHo3.1 is based on pGPTVKan (Becker, D. et al., Plant Molecular Biology 20 (1992), 1195-1197) with the addition of the 987bp gbss promoter cloned at the HindIII site of pGPTVKan and the uidA gene is deleted by Smal and SstI.

The mother variety Dinamo is transformed with the construct pHAbe12A as described in example 2.

10

Example 4

Down-regulation of StGH1 and StGH2 genes in potato by antisense

The StGH1 and StGH2 genes were down-regulated in potato by transformation with the genes in antisense direction in relation to a plant regulatory element. The respective antisense genes were cloned in a binary vector driven by a tuber specific gbss promoter. Nptll, yielding resistance to the antibiotic kanamycin, was used as selection marker. Two varieties were transformed, Prevalent and Producent. The shoots were selected on 50 μM kanamycin, which is a standard kanamycin concentration used for potato transformation (Ooms, G et al., Theoretical and Applied Genetics 73:744-750 (1987) and Tavazza, R. et al., Plant Science 59 (1988), 175-181).

Example 5

25

30

Over-expression of StGH1 and StGH2 genes in potato

The StGH1 and StGH2 genes were over-expressed in potato driven by the tuber specific promoter gbss. A mutated AHAS gene was used as selection marker yielding tolerance to the Imazamox herbicides. Two potato varieties were transformed, Desiree and AM99-2003 a transgenic high amylose line with a 40% decrease in starch content compared to its mother variety.

Example 6

35

Selection of transgenic lines

Non-transgenic escapes were identified and discarded by a PCR screening method.

DNA was extracted according to DNeasy 96 Plant protocol (Qiagen). In a 96 well microtiter plate, 10-15 mg leaf tissue was added to each well-together with a 5mm steel

ball, each well then representing one individual shoot. The plates were frozen in $N_2(1)$ before homogenisation. The homogenisation was done at 30Hz in a Mixermill300 for 1 min. The DNA was at the end of the extraction protocol eluted in 75 μ I H₂O.

5 Specific primers for nptII and AHAS were used for the amplification of a 246bp fragment respective a fragment of 509 bp for selection of successfully transformed lines.

Npt2_for 5'-AGCAAGGTGAGATGACAGGAGATC-3'
Npt2_rev 5'CAGACAATCGGCTGCTCTGATG-3'

10

AHAS1_frw: 5'-AACAACAACATCTTCTTCGATC-3' AHAS1_rev: 5'-TAACGAGATTTGTAGCTCCG-3'.

The PCR reactions were with the extracted DNA setup and run as follow:

15

Reaction:

10x PCR Mix 2,0 μl
Primer frw (25μM) 0,4 μl
20 Primer rev (25μM) 0,4 μl
dNTPs (10mM) 0,4 μl
RedTAQ (Sigma) 1,0 μl
Templat (~20ng/μl) 4,0 μl
H₂O 11,8 μl

25

30

PCR program:

94°C 30 s 59°C 30 s x29 cycles 72°C 30s 72°C 7 min 8°C Hold

Example 7

35

40

Gene expression analyses

The gene expression levels of the StGH1 and StGH2 genes were analysed in the transgenic potato lines with real-time PCR (ABI prism 7900HT, Applied Biosystems). With real-time PCR the change of gene expression can be analysed regarding RNA

expression levels. For pHS1 and pHS2 transgenic lines, expression of both sense and antisense RNA of StGH1 and StGH2 was measured, while in pHS3 and pHS4 transgenic lines the change in StGH1 and StGH2 mRNA expression was analysed. The target for pHS1 and pHS2 is to reduce transcript levels of StGH1 and StGH2 respectively while the target for pHS3 and pHS4 is to increase transcript levels of the respective genes.

RNA was isolated from microtubers of the transgenic potato lines and mother varieties using Invisorb Spin Plant-RNA mini kit (Invitek). A reverse transcription reaction was made with 250 ng total RNA in 25µl total reaction volume using TaqMan reverse transcription reagents (Applied Biosystems). Separate and specific primers (see table 1) were designed and used for the reverse transcription reaction in order to be able to differentiate the endogenous expression from the antisense RNA expression of the respective genes.

15

10

5

StGH1 sense RNA	5'-TGAAGACAGCACAAACTGG-3'
StGH1 antisense RNA	5'-GTGAAAGTTTGAACGCACAC-3'
StGH2 sense RNA	5'-AGTGCCATAACATGCTTTCC-3'
StGH2 antisense RNA	5'-CACATTTCAGCTGTTGATGGA-3'

Table 1

5µl of the reverse transcription reaction was used in triplicate analyses together with specific sequence detection primers, TaqMan MGB probe (see table 2) and UMM mastermix (Applied Biosystems) and determined with real-time PCR.

StGH1

Forward Primer: TCGAGTCGCCACGTAGAACTC Reverse primer: GAAATGCGTATGCGACTATGATG

TaqMan probe: AGTCTCTCGGAGTTCC

StGH2

Forward primer: GGTGCTGATCCTCCAGTTCTCT Reverse primer: GTCCCTGAAGCATAACCAAGGT

TaqMan probe: TTCTGCACTACTTAGGCCT

Table 2

Down-regulation of the two genes resulted in a decrease in gene expression in transgenic lines compared to their mother varieties in the order of 50-95 %.

Over-expression of the two genes resulted in a 2-10 times increase in gene expression in transgenic lines compared to their mother varieties.

				Times increase or de-
				crease in gene expres-
Ref no	Construct	Variate	1.00	sion compared to mother
P01-041-84	pHS1	Variety	ΔCts	variety
P02-325-1		Producent	-1,14	-1,3
	pHS1	Producent	-2,03	-4,1
P02-325-9	pHS1	Producent	-1,47	-2,2
P02-325-11	pHS1	Producent	-1,25	-1,6
P02-325-15	pHS1	Producent	-2,52	-6,3
P02-325-25	pHS1	Producent	-2,0	-4
P02-325-27	pHS1	Producent	-1,64	-2,7
P02-325-33	pHS1	Producent	-1,59	-2,5
P02-325-34	pHS1	Producent	-1,52	-2,3
P02-325-63	pHS1	Producent	-1,53	-2,3
P02-300-37	pH\$2	Prevalent	-1,27	-1,6
P02-300-66	pHS2	Prevalent	-1,04	1,1
P02-300-71	pHS2	Prevalent	-1,13	-1,3
P02-300-73	pHS2	Prevalent	-1,1	-1,2
P02-300-80	pHS2	Prevalent	-2,12	-4.5
P02-300-127	pHS2	Prevalent	-1,67	-2,8
P02-300-140	PHS2	Prevalent	-3,96	-15,7
P02-303-31	pHS2	Prevalent	-1,16	-1,4
P02-303-64	pHS2	Prevalent	-1,15	-1,3
P02-305-54	pHS2	Prevalent	-1,33	-1,8
P02-320-24	pHS2	Prevalent	-1,03	-1.1
P02-307-4	pHS3	Desirée	1,82	3,3
P02-307-5	pHS3	Desirée	2,68	7,2
P02-307-12	pHS3	Desirée	2,67	7,1
P02-307-14	pHS3	Desirée	1,83	3,3
P02-307-15	pH\$3	Desirée	1,79	3,2
P02-307-33	pHS3	Desirée	3,21	10,3
P02-307-43	pHS3	Desirée	2,7	7,3
P02-307-51	pHS3	Desirée	2,73	7,5
P02-307-80	pHS3	Desirée	2,78	7,7
P02-307-87	pHS3	Desirée	1,02	1,1

Ref no	Construct	Variety	Δ Ct s	Times increase or de- crease in gene expres- sion compared to mother
P02-307-148	pHS3	Desirée	1,88	Variety
P02-309-63	pHS3	AM99-2003	1,64	3,5
P02-309-106	pHS3	AM99-2003	1,75	2,7
P02-311-59	pHS3	AM99-2003	1,17	3,1
P02-312-15	pHS4	AM99-2003	1,03	1,4
P02-313-21	pHS4	AM99-2003	1,03	1,1 2,4

Table 3: Gene expression analyses based on Real- Time PCR

Example 8

5

10

15

20

Dry matter analysis

Dry matter has been analysed on microtubers from transgenic lines transformed with pHS1, pHS2, pHS3 and pHS4 showing a down-regulation or over expression of the genes. Since starch normally contribute to more than 80% of the dry matter in potato tubers, an increase or decrease in starch content will affect also the dry matter content.

Two microtubers of each line were harvested when they had reached maturity. Dry matter was calculated for mature microtubers weighed before and after 72 hours drying at 60°C. For comparison microtubers from the varieties Dinamo, Desiree, Prevalent, Producent and P737 with starch contents between 13 and 28% (when grown in field) were used. The starch content of microtubers is not as high as starch content of field grown tubers. However dry matter content can readily be compared in microtubers and that value is well correlated to the determined starch content in field grown tubers. In table 4 the average dry matter for the different varieties, calculated on ten or more microtubers, is shown.

Variety	Starch content field grown tubers	Dry matter microtubers
Dinamo	13%	14,8
Desirée	16%	16,1
Producent	22%	
Prevalent	22%	19,2
P737		19,7
1707	28%	21,6

Table 4. Dry matter content of 5 varieties based on 10 or more microtubers

One of each pHS1 and pHS2 with confirmed decrease in gene-expression have been analyzed for dry matter so far. Those two have a decrease in dry matter of 7 and 11% compared to their mother varieties.

For the pHS3 lines 8 of 9 of the confirmed over-expressed lines show an increase of up to 36% in dry matter. See table 5.

Ref No	Construct	Variety	Dry matter in relation to mother
		}	variety (%)
41-84	pHS1	Producent	89 -
300-127	pHS2	Prevalent	93
300-140	PHS2	Prevalent	96
307-4	pHS3	Desirée	106
307-5	pHS3	Desirée	117
307-15	pHS3	Desirée	124
307-33	pHS3	Desirée	116
307-57	pHS3	Desirée	136
309-63	pHS3	AM99-2003	134
309-106	pHS3	AM99-2003	109
309-111	pHS3	AM99-2003	108

Table 5. Dry matter content on transgenic lines with confirmed down-regulation or over expression of the StGH1 and StGH2 genes

Example 9

15

Starch content analysis

For analysis of starch content a total starch assay procedure from Megazyme International Ireland Ltd., Bray, Co.Wicklow, Ireland (AOAC Method 996.1; AACC method 76.13; ICC standard method No. 168) was used according to the suppliers instructions.

20 Starch content was analysed on microtubers from all transgenic lines transformed with pHS1, pHS2, pHS3 and pHS4. The microtubers were harvested when they had reached maturity. Mature microtubers were ground and maltosaccharides and free glucose residues were washed away with ethanol. The microtuber starch was treated with DMSO to ensure the complete solubilisation of samples with high levels of resistant starch, as the high amylose clones.

Samples were analysed with a standard spectrophotometric assay procedure. The transgenic lines were compared to potato varieties with known starch content ranging from 8% to 30%. The results give an indication on the change in starch content related to the genetic modification of the different transgenic lines.

5

Example 10

Field-testing of transgenic potato lines

- Produced transgenic lines are tested in field trials for determination of agronomic performance in relation to mother variety and other varieties used for starch production. Starch content, which is a main agronomic factor of importance for crops used for starch processing, can measured by several different methods.
- Under water weighing is performed on a scale in a tub of water and where an increase in starch content is associated with an increase in density of the sample. An increase starch is associated with an increased dry matter content, which can be measured by comparing the tissue fresh weight to tissue dry weight after extensive water elimination in an oven.

20

Starch content can also be measured by enzymatic methods as described under starch content analyses.

Example 11

25

30

Increased amylose and starch yield in potatoes

Potato varieties used for starch production as well as genotypes with a high amylose content are transformed with gene constructs as described above for the over-expression of a starch biosynthesis enhancing protein. The over-expression of StGH1 or StGH2 in potato plants will result in an increased armylose content of the transgenic plant compared to the starting plant.

Example 12

35

increased solids and improved processing quality of potatoes

In another aspect the invention may be used to increase the solids content of varieties that are used for processed potato products or table potato varieties. The potato genotypes are transformed with gene constructs as described above for the over-expression

of a starch biosynthesis enhancing protein. This starch biosynthesis enhancing protein may be derived from genes described above or other plant genes containing the same functional domains.

What is claimed:

5

- A method of increasing the production of starch in plants comprising culturing a
 plant with enhanced expression or activity of at least one starch biosynthesis
 enhancing protein.
- The method of claim 1, wherein said starch has a high amylose content.
- 3. The method as claimed in either claim 1 or 2, wherein production of amylose is increased.
 - 4. The method as claimed in any one of claims 1 to 3, wherein said method comprises over-expression of an amylose biosynthesis enhancing protein.
- The method as claimed in claim 4, wherein said protein comprises the SEQ ID NO: 2 or 4 or a protein derived from this sequence by substitution, insertion or deletion of amino acids and which has at least 50% identity at the amino acid level with SEQ ID NO: 2 or 4.
- The method as claimed in any of claims 1 to 5, wherein the amylose biosynthesis enhancing protein is encoded by a nucleic acid sequence selected from the group consisting of:
- a) a nucleic acid sequence comprising a nucleotide sequence which is at least 60% identical to the nucleic acid sequence of SEQ ID NO: 1 or 3;
 - a nucleic acid sequence comprising a fragment of at least 30 nucleotides of a nucleic acid sequence comprising the nucleotide sequence of SEQ ID NO:1 or 3;
 - a nucleic acid sequence which encodes a polypeptide comprising an amino acid sequence at least about 60% identical to the amino acid sequence of SEQ ID NO:2 or 4 and
- d) a nucleic acid sequence which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO: 2 or 4 or wherein the fragment comprises at least 10 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2 or 4,

15

20

35

- 7. The method as claimed in any one of claims 1 to 6, wherein the amylose biosynthesis enhancing protein is encoded by a nucleic acid sequence comprising the nucleotide sequence set forth as SEQ ID NO:1 or SEQ ID NO:3.
- 5 8. The method as claimed in any one of claims 1 to 7, wherein deficiency or decreased activity is achieved by a method selected from the group consisting of:
 - a) knock-out of the gene encoding said protein;
 - b) mutagenesis of the gene encoding said protein, wherein said mutation can be induced in the coding, non-coding, or regulatory regions of said gene;
 - c) expression of an anti-sense RNA, wherein said anti-sense RNA is complementary to at least part of the RNA encoding said protein;
 - 9. A method of producing amylose type starch by culturing a plant which overexpresses SEQ ID NO:1 or 3 or has increased amylose biosynthesis enhancing activity under conditions such that the plant produces an increased amount of amylose type starch.
 - 10. The method of any of the preceeding claims, wherein said plant belongs to the genus Solanum.
- 25 11. The method of claim 10, wherein said plant is Solanum tuberosum.
 - 12. A nucleic acid sequence SEQ ID NO:1 encoding an amylose blosynthesis enhancing protein.
- 30 13. A nucleic acid sequence SEQ ID NO:3 encoding an amylose biosynthesis enhancing protein.
 - 14. An amino acid sequence SEQ ID NO:2 having amylose biosynthesis enhancing activity.
 - 15. An amino acid sequence SEQ ID NO:4 having amylose biosynthesis enhancing acitivity.

10

15

20

30

- 16. A transgenic expression cassette comprising in combination with a regulatory sequence a nucleic acid sequence selected from the group consisting of:
 - a nucleic acid sequence comprising a nucleotide sequence which is at least 60% identical to the nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3,
 - a nucleic acid sequence comprising a fragment of at least 30 nucleotides of a nucleic acid sequence comprising the nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3,
- a nucleic acid sequence which encodes a polypeptide comprising an amino acid sequence at least about 60% identical to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4, or
 - a nucleic acid sequence which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4 wherein the fragment comprises at least 10 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4

wherein said regulatory sequence is capable of mediating expression of said nucleic acid sequence in a plant.

- 17. A transgenic expression cassette of claim 16, wherein said regulatory sequence is a promoter sequence heterologous with regard to said nucleic acid sequence.
- 18. A transgenic expression cassette of claim 16, wherein said regulatory sequence is a tuber specific promoter sequence.
 - 19. A transgenic expression cassette of either claim 16, 17 or 18, wherein said nucleic acid sequence is arranged in antisense or sense orientation with regard to said promoter sequence.
 - 20. A transgenic expression cassette of any of the claims 16 to 19, wherein said nucleic acid sequence encodes a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:4.
- 35 21. A transgenic expression cassette of any of the claims 16 to 20, wherein said nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO:1 or SEQ ID NO:3.

15

- 22. A transgenic expression cassette of any of the claims 16 to 21, wherein said nucleic acid sequence encodes a naturally occurring variant of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:4:
- 5 23. A transgenic host cell transformed with an expression cassette of any of the claims 16 to 22.
 - 24. A transgenic host cell of claim 23, wherein said host cell belongs to the genus Solanum.
 - 25. A transgenic plant comprising an expression cassette of any of claims 16 to 22.
 - 26. A transgenic potato plant comprising an expression cassette of any of claims 16 to 22.
 - 27. A transgenic potato plant, plant part, seed or tuber comprising an expression cassette of any of claims 16 to 22.

SEQUENCE LISTING

	<110> BASF Plant Science GmbH
	5 <120> Enhanced Amylose Production in Plants
	<130> AE884-02
1	<140> PF0000054331
,	
	<160> 15
15	<170> PatentIn Ver. 2.1
	<210> 1
	<211> 2084 <212> DNA
00	<213> Solanum tubomasi
20	<220>
	<221> CDS
	<222> (302)(1696)
25	<400> 1
	ttttcataaa cttcttcaac tttattccat actcttttat ttatcagctc ctagatcttc 60
	ttttttgttt gttgatatte tetraageet totageet tetraageet totagetete 60
30	ttttttgttt gttgatattc tcttgaaaat tgttcagtga agagttgatc aaagctaaga 120
	cacagggget geggecattt ttteacegga atettettet ttattteeeg gtgaaagttt 180
	gaacgoacac cgttatttct agacagtaga caatgtcaag tgaaaaacat cacaágtttt 240
35	tgaagatttg taattaatta gttgagattt ttaatttgga ggaaagagaa aaacagagaa 300
	g atg ata ggg cgg gtg ggg the the the
	"
40	10 15
40	gtg act att ggg gct gaa acg acg acg tta aaa ggg gta aac aga aat 397
	20 20 Asn Arg Asn
	30
45	gcg tat gcg act atg atg tat atg gga act ccg aga gac tac gag ttc 445 Ala Tyr Ala Thr Met Met Tyr Met Gly Thr Pro Arg Asp Tyr Glu Phe
	35 40 Asp Tyr Glu Phe
	tac gtg gcg act cga gta atg ctc cga tca ctt acc cgg cta gga gtt 493
50	The same and the per their the law that care and
	50 55 60
	gaa gcc gat ctc gtc gtt att gct tca ctt gac gtt cct ctc egc tgg 541
	65 TO Leu Arg Trp
55	,2 80
	gtt caa act cta gaa cag gaa gat ggt gct aag gtg gtg aga gtt aaa 589
	Val Gln Thr Leu Glu Gln Glu Asp Gly Ala Lys Val Val Arg Val Lys 85 90 95
60	
	aat ctg aac aat ccg tat tgt atc aac cct aat tgg aga ttc aag ctc 637 Asn Leu Asn Asn Pro Tyr Cys-Ile Asn Pro Asn Trp Arg Phe Lys-Leu
	100 105 110 ASI TIP ARG PRE LYB LEW -

	aca ctg aac aaa ctt tat gcg tgg agc ctc gta aat tat gac agg gtt 685 Thr Leu Asn Lys Leu Tyr Ala Trp Ser Leu Val Asn Tyr Asp Arg Val 125
•	5 gtc atg ctt gat gct gac aac ctt ttc ctc cag aaa act gat gaa ctg 733 Val Met Leu Asp Ala Asp Asn Leu Phe Leu Gln Lys Thr. Asp Glu Leu 130
1	ttc caa tgt ggc cag ttt tgt gct gtc ttc att aat ccc tgc atc ttc 781 Phe Gln Cys Gly Gln Phe Cys Ala Val Phe Ile Asn Pro Cys Ile Phe 150 150 160
19	170 175 ASD 176
20	CBA dot tit att and and
25	Caa ggt tit att gga ggc cac ttc cca gat tta cit gat cgg cca atg 925 Gln Gly Phe Ile Gly Gly His Phe Pro Asp Leu Leu Asp Arg Pro Met 200 205
	Phe His Pro Pro Leu Asn Gly Thr Gln Leu Gln Gly Ser Tyr Arg Leu 210 215 220
30	Cct cta gga tac caa atg gac gcc tct tat tat tat ctc aaa ctc cat 1021 Pro Leu Gly Tyr Gln Met Asp Ala ser Tyr Tyr Tyr Leu Lys Leu His 235 240
35	tgg tcg gta cct tgt gga cct aat agt gtc att aca ttt cct ggt gct 1069 Trp Ser Val Pro Cys Gly Pro Asn Ser Val Ile Thr Phe Pro Gly Ala 245 250 255
40	Pro Trp Leu Lys Pro Trp Tyr Trp Ser Trp Pro Val Leu Pro Leu 260 265 270
45	ggc atc cag tgg cat gaa cag cga cgt cta act gtt ggg tat ggt gct 1165 Gly Ile Gln Trp His Glu Gln Arg Arg Leu Thr Val Gly Tyr Gly Ala 285
70	gag atg ata gca gtg ttg atc caa tct ata ttt tac cta gga ata att Glu Met Ile Ala Val Leu Ile Gln Ser Ile Phe Tyr Leu Gly Ile Ile 290 295 300
50	gca gtg aca cgc cta gca cgc cca aat tta tca aag ttg tgc tat cgc 1261 Ala Val Thr Arg Leu Ala Arg Pro Asn Leu Ser Lys Leu Cys Tyr Arg 315 320
55	Cat gat agc aag agt gcc ttc tta cta cga act ggc ctt aaa ttg 1309 Ris Asp Asp Ser Lys Ser Ala Phe Leu Leu Arg Thr Gly Leu Lys Leu 335
60	att gct ata tgg tcc att ctt gct gcc tac aca gtt cct tat ttc gtg 1357 Ile Ala Ile Trp Ser Ile Leu Ala Ala Tyr Thr Val Pro Tyr Phe Val 340 345 350
	att cct tgt aca gtt cat cca cta gtt ggc tgg agt ctc tac tta ctc 1405 Ile Pro Cys Thr Val His Pro Leu Val Gly Trp Ser Leu Tyr Leu Leu

	355	. 360	365	
5	ggc tct ttt tca cta tcc Gly Ser Phe Ser Leu Ser 370	tgt ata aca Cys Ile Thr 375	gtg aat gca ttt ctt ttg ccg Val Asn Ala Phe Leu Leu Pro 380	145
10	385 390	. Tro tro tre	ggg atc ctt ggg gcc ctt ttg Gly Ile Leu Gly Ala Leu Leu 395 400	1501
	gtg atg gct tac cct tgg Val Met Ala Tyr Pro Trp 405	-3- ver veh	ggt gtt gta aga gca atg gct Gly Val Val Arg Ala Met Ala 415	1549
15	gta tit aca tac gcc ttc Val Phe Thr Tyr Ala Phe 420	tgt get tet Cys Ala Ser 425	cca gca tta tgg atg gca ttg Pro Ala Leu Trp Met Ala Leu 430	1597
20	gtt aaa atc aag tgt tct Val Lys Ile Lys Cys Ser 435	ctt cat gtt : Leu His Val :	tca ctt gag agg gaa gga ttc Ser Leu Glu Arg Glu Gly Phe 445	1645 [°]
25	ttg ccc aag ata agt gaa Leu Pro Lys Ile Ser Glu 450	tct aca gca c Ser Thr Ala E 455	ect get ggt tet aac aaa etg Fro Ala Gly ser Asn Lys Leu 460	1693
30	tat tgaaagttga aaagttaaa Tyr 465	g gaatcaacag	gagaactaat gcttcagaaa	1746
O	catctccaaa cgttttgctt ag	gagacttg gagt	ctgctt gtgctatcct agctagttgc	1806
			tgagtg ggtttagaat tgggaggatg	
35			gaatgo tactccagga aaaagattgt	
			gtggaa taggttettt aaaatttatt	
40			tattaa gtootggtga aataagtaat	
	toottattoa tacaaaaaaa aaa			2084
45	<210> 2 <211> 465 <212> PRT <213> Solamum tuberosum		,	
50	<400> 2 Met Ile Gly Arg Val Gly L		d Leu Leu Ile Ala Thr Thr 0 15	
55	Val Thr Ile Gly Ala Glu T 20	hr Thr Thr Le 25	u Lys Gly Val Asn Arg Asn 30	•
	Ala Tyr Ala Thr Met Met T 35	yr Met Gly Th 40	r Pro Arg Asp Tyr Glu Phe 45	
60	Tyr Val Ala Thr Arg Val M	et Leu Arg Se 55	r Leu Thr Arg Leu Gly Val	
	Glu Ala Asp Leu Val Val I	le Ala Ser Le	u Asp Val Pro Leu Arg Trp	

		65			:.	ī		70						75							80
5	v	al (ln	Thz	Le	u G	lu G 3.5	ln G	lu	Asj) G]	y A	la : 90	Lys	٧a	1 V a	al <i>8</i>	ırg	Va 9		Lys
·		sn I	eu i	Asn	10	n P1 0	o T	YX C	λŧ	Ile	As 10	n P 5	ro 1	Asn	Txj	o Aı		he 10	Lу	s I	ieu
10	Tì	ır L	eu 2	Asn L15	Ly	s Le	u Ty	Æ A	la	Trp 120	Se	r L	eu T	7al	Ası	1 Ty	r A	ga	Arg	v	al
	Va	1 M	et 1 30	Leu	Ası	Al	a As	р Аз 13	511 : 35	Leu	Ph	e Le	<i>en</i> €	ln	Lys 140	Th	r A	sp	Glu	ı L	eu
15	Ph 14	e G: 5	ln C	у́в	GT?	/ Gl:	n Ph 15	e Cy O	78 7	Ala	Va.	l pł	ie I 1	le SS	Asn	Pr	o C;	ys	Ile		be 60
20	Hi	s Ti	r G	ly	Leu	16!	≥ Va	l Le	u (31n	Pro	Se 17	E L	ys :	Lys	Va:	l Pi		Asn 175		sp
	Me	t II	e #	is	Glu 180	Ile	Gl:	ı Il	e G	Sly	Arg 185	Gl	u A	sn (3 ln	Asp	G] 19		Ala	Αε	φ
25	Gli	n Gl	y Pi 1:	he 95	Ile	Gly	Gly	/ Hi	s P 2	he 00	Pro	As	p Le	eu 1	Leu	Asç 205	Ar	g 1	Pro	Mę	ŧ
	Phe	21	s Pi O	ro :	Pro	Leu	. Agr	Gl; 21!	у Т 5	hr	Gln	Lei	1 G]	.n. 6	31y 20	Sex	ту	r A	rg	Le	ы
30	Pro 225) Le	u G]	y !	Tyr	Gln	Met 230	Ası	A	la .	Ser	Tyz	7 Ty	T I 5	'nΣ	Leu	Ly	ș I	⁄eu	Hi 24	
35	Txp	Se:	r Va	1.1	Pro	Сув 245	Gly	Pro	A	sn i	Ser	Va] 250	. Il	е т	hr	Pļhe	Pr		ly 55	Al	à
	Pro	Tr	Le	u I 2	660 FA8	Pro	Trp	Тух	T	rp !	1 10 265	Sez	Tr	p P	ro '	Val	եеւ 27(ro	Lei	ıı
40	•	•	-,	_			Glu		25	5 U						285					
							Leu	473						3(90			•			•
45	Ala 305	Val	Th	r A	rg :	Leu	Ala 310	Arg	Pr	o A	sn	ŗėń	Se:	r Ly	ys 1	Leu	СУ¤	T		Arg 320	
50	His	Asp	Asj	o S	er :	Lys 325	Ser	Ala	Ph	e L	eu	Leu 330	Arg	Tì	ır (ly	Leu	163 33		Leu	ı
	Ile	Ala	Ile	3 T.	rp : 40	Ser	Ile	Leu	Al	a A 3	la 45	Tyr	Thi	. Va	ıl E		Tyr 350	Ph	le 1	/al	•
55 .	Ile	Pro	Cys 355	s 17] S	br 1	/al	His	Pro	Lе 36	u V	al (Gly	Tre	S e		eu 65	Tyr	Le	u I	ieu	
	Gly	Ser 370	Phe	e Se	er 1	ceu	Ser	Сув 375	Il	e T	hr '	Val	As n	Al 38		he	Leu	Le	u I	ro	
60	Met 385	ren	Pxc	> Va	al I	eu '	Val 390	Pro	Tx	p I	le (31y	Ile 395	Le	uG	ly :	Ala	Le		eu 00	

Val Met Ala Tyr Pro Trp Tyr Asn Asp Gly Val Val Arg Ala Met Ala 410 Val Phe Thr Tyr Ala Phe Cys Ala Ser Pro Ala Leu Trp Met Ala Leu 5 Val Lys Ile Lys Cys Ser Leu His Val Ser Leu Glu Arg Glu Gly Phe 440 10 Leu Pro Lys Ile Ser Glu Ser Thr Ala Pro Ala Gly Ser Asn Lys Leu Tyr 465 15 <210> 3 <211> 2230 20 <212> DNA <213> Solanum tuberosum <220> <221> CD\$ 25 <222> (143)..(2086) tatccccaga gaatcagctg aatcaagaac tgatttttag attatgtttt cttgattctt 60 30 tgaaatggga acttgatttt cagtttttca actcagatgt tgtgttcctt tagctggaaa 120 acttgaaaaa ggaaagccca ga atg aga gga agt tta got ggt gga cca cct Met Arg Gly Ser Leu Ala Gly Gly Pro Pro 35 agt cot att gaa cot aga cag agg ctt tot gta tto act gag gaa aca Ser Pro Ile Glu Pro Arg Gln Arg Leu Ser Val Phe Thr Glu Glu Thr 220 40 age aaa aga agg tto ttg aga agt aaa gtt tte aga gat ggg gag aga Ser Lys Arg Arg Phe Leu Arg Ser Lys Val Phe Arg Asp Gly Glu Arg 268 get ett cat agt eec ace aaa aac agg aat tit ace tge aag tie eea 45 Ala Leu His Ser Pro Thr Lys Asn Arg Asn Phe Thr Cys Lys Phe Pro 316 act gtg aag ctt ata ttg ggt gtt att gct ctg gtt gca att tgg tca Thr Val Lys Leu Ile Leu Gly Val Ile Ala Leu Val Ala Ile Trp Ser 364 50 ctc tgg cat tct cca gca att tat aac acg gaa tac ata tct agt tca Leu Trp His Ser Pro Ala Ile Tyr Asn Thr Glu Tyr Ile Ser Ser Ser 55 ggc tot cgg get get ttg atg cac aga gag tta agt ggt cat tet tea Gly Ser Arg Ala Ala Leu Met His Arg Glu Leu Ser Gly His Ser Ser 95 get gat caa egt tat aca tea ett tta gat att gae tgg gae caa att Ala Asp Gln Arg Tyr Thr Ser Leu Leu Asp Tle Asp Trp Asp Gln Ile 120

. 5		er G	aa g ln V 1	rtt a Val I. 25	tt g le G	ag aa lu Ly	a ci	. u . A,	cc g la A 30	at a sp A	gg c	at g is G	lu T	at c yr G 35	ag g	gc gt: ly Va	a 556 L
	G)	-	ta t le L 40	ta aa eu Aa	ac ti en Pl	tc aa ne As	t ga n As 14	.p 36	gt ga er Gl	e al lu II	it gr le Ar	Bp G	ag t ln L	tg a eu L	ag g Ys G	ag tta lu Ler	604
10		a co u Pi 5	eg ga	ac go ep Al	t ga .a Gl	g ca u Hi	5 V 64	a at l Il	c tt	g as u As	ic ct in Le 16	u As	at ca sp Hi	ac gt is Va	tc co	ng aat ro Asn 170	
15	aa As:	t at n Il	a ac .e Ti	ia tg ir Tr	g ga p Gl 17		a at	a ta e Ty	t do r Pr	t ga o Gl 18	u TT	g at p Il	a ga e As	ıt ga Ip Gl	ia ga .u G]	aa gaa lu Glu IS	700
20				19	0	- 4111	. Cya	3 FL(19	r Lei	u Pr	o Ly	s Il	e Gl 20	n Ph O	t ccg e Pro	748
25	Ī	•	20	5		, mp	- 1	210)	. val	ı ryı	9 Let	21:	D Су: 5	s Ly	a Lya	796
		220		<u>-</u> -	, - ,,	<u></u>	225	AGT	Ala	AIÇ	, bπ€	230	3 Let)	ı Glı	1 Lei	g gca u Ala	844
30	235		- •			240	CCL	ASH	пyв	сту	245	HIS	Pro) Ile	His	t gtg s Val 250	892
35					255	. ALS	ъпе	1. LO	THE	260	Asn	Leu	Phe	Thr	265		940
40				270	AL 9	GIU	GIŸ	ASD	275	TXD	Leu	Tyr	Glu	Pro 280	Asn	ctg Leu	988
45			285	****	Gia	пур	Leu	290	ren	Pro	val	Gly	Ser 295	Cys	Glu	ctt Leu	1036
		300		204	шув		19 8 305	ATG	ASN	Trp	His	Ser 310	Gly	Asn	Val	Arg	1084
50	cga Arg 315	gaa Glu	gcc Ala	tat Tyr	gca Ala	act : Thr : 320	att Ile	ctc Leu	cac His	tca Ser	gca Ala 325	aat Asn	ttt Phe	tat Tyr	gta Val	tgt Cys 330	1132
55	gga Gly	gec Ala	ata Ile	wra	gca Ala: 335	gca (Ala (cag Sln	agt Ser	IT6	cgc Arg 340	ttg Leu	gca Ala	ggt Gly	tca Ser	acc Thr 345	cga Arg	1180
- 60	gat (Asp	ctt Leu	gtg Val	ata Ile 350	ctt Leu	gtt (Val)	yat Asp	GIU	act Thr 355	atc Ile	agt Ser	gac Asp	tac Tyr	cac His 360	agg Arg	ggt Gly	1228
	ggt 1	tta	gag	gct	gcc ·	gga t	-99 ·	aag	atc (cac	acg	ata	aag	aga	ata	agg	1276

	43		•			/			
				3	70		le Lys Arg 375		
5	380	•		385	TO TAT 7	35 ABH GIU T	gg aac tat rp Asn Tyx 90	Ser Lys	1324
10	395	-	400)	eb iår t	405	tc atc ttc le Ile Phe	Ile Asp 410	1372 _.
15	•		415	. wra we	A TIE	asp Phe Le	co ttt gag eu Phe Glu	Met Pro 425	1420
20		430	dec dry	WOTH WE	435	er ren by	t aat tca e Asn Ser 440	Gly Val	1468
20		445	-140 501,	45	o The P.	ue GIN l'é	g ttg.atg u Leu Met 455	Asp His	1516
25	460		002	465	n Giy G.	LY ASD GIN	•	Leu Asn	1564
30	475		480	WIS WIG	1 TTG 152	CO Lys His 485	e atg aac s Met Aşn	Phe Leu 490	1612
35	,	7	495	web are	50	o Tring Lys		Lys Thr. 505	1660
		510	ara veh	rio Pio	515	u Tyr Val	ctg cac t Leu His : 520	Tyr Leu	1708
40		aa cct ys Pro : 25	tgg tta Trp Leu	tge tte Cys Phe 530	Arg As	c tac gat p Tyr Asp	tgc asc t Cys Asn 1 535	igg aat Irp Asn	1756
45	gtg ggt a Val Gly L 540	ag ttg (ys Leu (JEN GIG	ttt gca Phe Ala 545	agt gar Ser Asj	t gtg gca p Val Ala 550	cac agg a His Arg T	icg tgg Thr Trp	1804
50	tgg aag gu Trp Lys Va 555		560	det bio	ASP ASI	n Leu His 565	Lys Tyr C	570 570	1852
5 5	ctt agg to Leu Arg Se	5	75	ra wia	580	Trp Asp	Arg Arg G	lu Ala 85	1900
	gag aaa go Glu Lys Al	590	502 2	ab GIÀ	595	h TAR ITE	eoo ra 116 ri	ys Asp	1948
60	pro Arg Le		ct tgt t br Cys T	at gaa Yr Glu 610	gaa ttt Glu Phe	Cys Phe	tgg gaa aq Trp Glu se 615	gc atg er Met	1996

5	ŗe	u Tr 62	b wr	ic tg .s Tr	b er	r ga y Gl	a ac u Th 62	r As	c tg n Tr	g ac p Th	a ga r As	t aa p As 63	n Al	c ac a Th	c tc x Se	t tca r Ser	2044
	ec Pro 63!	o Th	a co r Pr	t cc o Pr	c at o Mei	g gto t Val 640	L As:	t ac	t gçi r Ala	t tca a Sei	a cti r Lei 64!	u Se	t tc r Se	t tt	n a		2086
10	taa	acta	ggag	gtt	ctgti	tac t	atac	ctc	gc ta	gtict	gtaa	a gt	ata	caga	gte	acaact	t 2146
	gaa	tgt	caac	cag	tgcca	att a	gtat	acat	g gt	gtca	acac	: tt	taca	eacc	ctts	gaaaaa	a 2206
15	aaa	laaa	aaaa	aaaa	aaaa	laa a	aaa										2230
20	<21 <21	.0> 4 .1> 6 .2> I .3> 5	548 PRT	um t	uber	Osum	l					·				٠	
25	Met 1		Gly		5					10					1.5		
	Gln	Arg	Leu	Ser 20	Val	Phe	Thr	Glu	Glu 25	Thr	Ser	Lys	Arg	Arg 30		Leu	
30	Arg	ser	Lys 35	·Val	Phe	Arg		Gly 40	Glu	Arg	Ala	Leu	His 45		Pro	Thr	
•	Lys	Asn 50	Arg	Asn	Phe	Thr	Сув 55	Lys	Phe	Pro	Thr	Val 60	Lys	Leu	Ile	Leu	
35	Gly 65	Val	Ile	Ala	Leu	Val 70	Ala	Ile	Ţŗp	Ser	Leu 75	Trp	His	Ser	Pro	Ala 80	
40	Ile	Tyr	Asn	Thr	Glu 85	Tyr	Ile	Ser	Ser	Ser 90	Gly	Ser	Arg	Ala	Ala 95	Leu	
	Met	His	¥rg	Glu 100	Leu	Ser	Gly	His	Ser 105	Ser	Ala	Asp	Gln	Arg 110	Tyr	Thr	
45	Ser	Leu	Leu 115	Asp	Ile	qaA	Trp	Asp 120	Gln	Ile	Ser	Gln	V al 125	Ile	Glu	Lys	
	Leu	Ala 130	Asp	Arg	His	Glu	тут 135	Gln	Gly	Val	Gly	Ile 140	Leu	Asn	Phe	Asn	
50	Asp 145	Ser	Glu	Ile	Asp	Gln 150	Leu	īys	Glu	Leu	Leu 155	Pro	.Asp	Ala	Glu	His 160	:
EE	Val	Ile	Leu	Asn	Leu 165	Asp	His	Val	Pro	Asn 170	Asn	Ile	Thr	Trp	Glu 175	Thr	
55	Ile	Tyr	Pro	Glu 180	Trp	Ile	Asp	Glu	Glu 185	Glu	Glu	Phe	Glu	Val 190	Pro	Thr	
60	Cys	Pro	Ser 195	Leu	Pro	Lys	Île	Gln 200	Phe	Pro	Gly	Lys	Pro 205	Arg	Ile	Asp	
	Leu	īle	Val.	Val	Lys	Leu	Pro	Cys	Lys	Lys	Sex	Lys	Aso	Tro	Tvr	Ara	

35

50

	210	ı				215			•		220	٠				
5	Asp Val 225	Ala	Arg	bpė	His 230	Leu	Gln	Leu	Ala	Ala 235	Ala	Arg	Leu	Äla	Ala 240	
	Ser Asn	Lys	Gly	Tyr	His	Pro	Ile	nis	Val	Leu	Leu	val	Thr	Glu	His	

245

Phe Pro Thr Pro Asn Leu Phe Thr Cys Lys Glu Leu Val Val Arg Glu 10 265

Gly Asn Ala Trp Leu Tyr Glu Pro Asn Leu Asn Thr Leu Arg Glu Lys

Leu His Leu Pro Val Gly Ser Cys Glu Leu Ala Val Pro Leu Lys Ala 15

Lys Ala Asn Trp His Ser Gly Asn Val Arg Arg Glu Ala Tyr Ala Thr 310

Ile Leu His Ser Ala Asn Phe Tyr Val Cys Gly Ala Ile Ala Ala Ala 330

Gln Ser Ile Arg Leu Ala Gly Ser Thr Arg Asp Leu Val Ile Leu Val 25

Asp Glu Thr Ile Ser Asp Tyr His Arg Gly Gly Leu Glu Ala Ala Gly 355

30 Trp Lys Ile His Thr Ile Lys Arg Ile Arg Asn Pro Lys Ala Glu Gln

Asp Ala Tyr Asn Glu Trp Asn Tyr Ser Lys Phe Arg Leu Trp Gln Leu 395

Thr Asp Tyr Asp Lys Ile Ile Phe Ile Asp Ala Asp Leu Leu Ile Leu

Arg Asn Ile Asp Phe Leu Phe Glu Met Pro Glu Ile Thr Ala Ile Gly 40

Asn Asn Ala Thr Leu Phe Asn Ser Gly Val Met Val Val Glu Pro Ser 440

Asn Cys Thr Phe Gln Leu Leu Met Asp His Ile Asn Glu Ile Glu Ser 45

Tyr Asn Gly Gly Asp Gln Gly Tyr Leu Asn Glu Ile Phe Thr Trp Trp 465 470

His Arg Ile Pro Lys His Met Asn Phe Leu Lys His Tyr Trp Glu Gly

Asp Glu Glu Lys Lys Gln Met Lys Thr Arg Leu Phe Gly Ala Asp 55 505

Pro Pro Val Leu Tyr Val Leu His Tyr Leu Gly Leu Lys Pro Txp Leu 520

60 Cys Phe Arg Asp Tyr Asp Cys Asn Trp Asn Val Gly Lys Leu Gln Glu 530 535 540 ----

		•								. •							
	Phe 545	Ala	Ser	ýsb	v al	Ala 550	His	Arg	Thr	Trp	Trp 555	Lys	Val	His	Asp	Ala 560	
5	Met	Pro	qaA	Asp	Leu 565	His	Lys	тух	Сув	Leu 570	ren	Arg	Ser	ГУB	Gln 575	Lys	
	Ala	Ala	Leu	Glu 580	Trp	Asp	Arg	Arg	Glu 585	Ala	Glu	Lys	Ala	Asn 590	Phe	ser	
10	Asp	Gly	His 595	Trp	Lys	Ile	Lys	Ile 600	ГÀв	Ąsp	Pro	Arg	Leu 605	Glu	Thr	Сўв	
	Tyr	Glu 610	Glü	Phe	Cys	Phe	Trp 615	Glu	Ser	Met	Leu		His	Trp	Gly	Glu	
15	Thr 625	Asn	qxT	Thr	Asp	Asn 630	Ala	Thr	Ser		Pro 635	Thr	Pro	Pro	Met	Val 640	-
20	Asn	Thr	Ala	Ser	Leu 645	.Ser	Ser	Leu									
25	<210 <211 <212 <213	> 14 > DN	ra.	lopsi	is th	nalia	una									,	
30	<220 <221 <222	> CI		(1485	5)												
35	<400 atg Met 1	gat	tta Leu	caa Gln	aga Arg 5	act Thr	ttg Leu	atg Met	ttc Phe	tct ser 10	tgt Cys	tgg Trp	gtt Val	ctg Leu	tct Ser 15	ctt Leu	48
·40	ttg Leu	atc Ile	atc Ile	aaa Lys -20	acg Thr	aca Thr	gcg Ala	tat Tyr	aac Asn 25	gag Glu	aaa Lys	cag Gln	ctg Leu	ttc Phe 30	cag Gln	Pro CCG	96
45	ctt Leu	gaa Glu	acg Thr 35	gaa Glu	aac Asn	gca Ala	aac Asn	gcg Ala 40	atg Met	acc Thr	Ala gcg	gtt Val	atg Met 45	gag Glu	cga Arg	gga gga	144
45	tta Leu	aag Lys 50	acg Thr	cag Gln	Arg	Arg	ccg Pro 55	gag Glu	cac His	aag Lys	aac Asn	gct Ala 60	tat Tyr	gcg Ala	acg Thr	atg Met	192
50	atg Met 65	tac Tyr	atg Met	gga Gly	aca Thr	cca Pro 70	aga Arg	gac Asp	tac Tyr	gag Glu	ttc Phe 75	tac Tyr	gtt Val	gcg Ala	aca Thr	cgt Arg . 80	240
55	gtc Val	ttg Leu	atc Ile	aga Arg	tcg Ser 85	ctt Leu	rås aag	agt Ser	ctc Leu	cac His 90	gtg Val	gac	gct Ala	gat Asp	atc Ile 95	gtc Val	288
60	gtt Val	ata Ile	gcc Ala	tcc Ser 100	Leu	gac Asp	gtt Val	cct Pro	atc Ile 105	Asn	tgg Trp	att Ile	cac His	gct Ala 110	Leu	gaa Glu	336
												aat	a++	~ ~ ~	aat	669	384

	Glı	a Glu	Asr 115	Gly	Ala	Lys	val	l Vaj	L Arg	y Va.	l Gl	u Ası	n Le 12		u As	n Pro	
5	tac Tyr	aag Lys 130	nys	caa Gln	acc Thr	aac Asn	Phe 135	: Asy	aac Asi	aga A Arg	tto J Phe	aaq Lyg 140	Le	t ag u Se:	t ct: r Lei	a aac u Asn	432
10	aag Lys 145	TIEN	tac Tyr	gct Ala	tgg Trp	tct Ser 150	ctic Leu	tct Ser	gat Asp	tat Tyr	gac Asp 155	Arg	get Val	gta l Val	a ato L Met	g ctt Leu 160	480
15	иц	VAL	nsp	watt	165	Pne	nen	Lys	Asn	170	Asp	Glu	Leu	Phe	175		528
	ej A aac	caa Gln	ttt Phe	tgt Cys 180	gct Ala	gtc Val	ttc Phe	atc	aac Asn 185	Pro	tgc Cys	atc Ile	ttc Phe	cac His	Thr	ggt	576
20	ctc Leu	ttt Phe	gtg Val 195	ttg Leu	Glu	cca Pro	tca Ser	atg Met 200	gag Glu	gtc Val	ttt Phe	aga Arg	gac Asp 205	atg Met	ctt Leu	cat His	624
25	gag Glu	ctt Leu 210	gaa Glu	gta Val	aag Lys	aga Arg	gat Asp 215	aac Asn	cet Pro	gat Asp	gga Gly	gct Ala 220	yab	caa Gln	cly ggc	ttt Phe	672
30	ctt Leu 225	gtc Val	agc Ser	tac Tyr	ttc Phe	tct Ser 230	gat Asp	tta Leu	ctc Leu	aat Asn	cag Gln 235	cct Pro	ata Leu	ttt Phe	cgt Arg	cct Pro 240	720
35	cct Pro	ccc Pro	gat Asp	ABI	cgc Arg 245	acc Thr	gcg Ala	ctt Leu	aag Lys	gga Gly 250	cat His	ttt Phe	agg A r g	ctt Leu	cct Pro 255	ttg Leu	768
	gga Gly	tat Tyr	GIN	atg Met : 260	gac Asp	gca Ala :	tct Ser	tat Tyr	tac Tyr 265	tac Tyr	ctt Leu	aag Lys	ctc Leu	aga Arg 270	tgg Trp	aac Asn	816
40	gta Val	cca Pro	tgt Cys 275	gga (cca :	aac : Asn !	ser	gtg Val 280	ata Ile	acg Thr	ttc Phe	cca Pro	gga Gly 285	gca Ala	gta Val	tgg Trp	864
45	Hen	290 Lys 38g	cca Pro	tgg (Trp (tat (Tyr !	rxb :	tgg Trp 295	tca Ser	tgg Trp	eet Pro	Val	ctt Leu 300	cct Pro	tta Leu	gly ggc	ctt Leu	912
50	tca ser 305	tgg Trp	cac (His)	cac (His (Gln 2	ego (Arg 1 310	ra Ge	tac Tyr '	acg Thr	Ile	agt Ser 315	tat Tyr	tca Ser	gca Ala	gag Glu	atg Met 320	960
55	cct Pro	tgg : Trp '	gtc (Val 1	Leu :	acc o Thr (caa q Fln A	gca Lla	gtg Val	Phe	tac Tyr 330	cta Leu	gga Gly	att Ile	ata Ile	cta Leu 335	gtc Val	1008
~~	aca .	cgt (Arg)	ren 1	gcg (Ala) 340	egt (Arg 1	ecc a Pro I	lac lsn	Met '	acc Thr 345	aag Lys	cta Leu	tgt Cys	Tyr	cga Arg 350	cgt Arg	tct \$er	1056
60	gat a	raa y	aat (Asn 1 355	Leu S	age a Ser N	atg a let]	(le (cag Gln (360	aca Thr	gct Ala	ttc Phe	ŗĀē :	ttt Phe 365	gtt Val	gca Ala	ctc Leu	1104

5	ctc Leu	Phe 370	atc Ile	ctc Leu	tca Ser	gcc Ala	tac Tyr 375	att Ile	ata Ile	cca Pro	ttc Phe	ttc Phe 380	atc Ile	atc Ile	cca Pro	cag Gln	1152
•	acg Thr 385	atc Ile	cac His	Pro ccg	ctc Leu	att Ile 390	ggt Gly	tgg Trp	tat Ser	ctc Leu	tac Tyr 395	tta Leu	acc Thr	ej ge	tcc ser	ttt Phe 400	1200
10	gct Ala	ctc Leu	tct Ser	acc Thr	ata Ile 405	ccc Pro	atc Ile	aac Asn	gcc Ala	ttc Phe 410	ttg Leu	ctt Leu	cc¢ Pro	att Ile	ctc Leu 415	ect Pro	1248
15	gtc Val	ata Ile	aca Thr	ccg Pro 420	tgg Trp	ctt Leu	Gly gg¢	att Ile	ttc Phe 425	gly 999	aca Thr	ctc Leu	ct¢ Leu	gtg Val 430	atg Met	gct Ala	1296
20	ttt Phe	ect Pro	tct Ser 435	tat Tyr	cct Pro	gat Asp	ggc	gtt Val 440	gtt Val	aga Arg	gca Ala	ttg Leu	tcg Ser 445	gtt Val	ttc Phe	eja aaa	1344
25	tat Tyr	gca Ala 450	ttt Phe	tgt Cys	tgt Cys	gca Ala	ccg Pro 455	ttt Phe	cta Leu	tgg Trp	gtc Val	tcc Ser 460	ttt Phe	gtg Val	aag Lys	atc Ile	1392
20	aca Thr 465	tcg ser	cat His	ctt Leu	cag Gin	att Ile 470	atg Met	att Ile	yeb	aaa Lys	gag Glu 475	gtt Val	ttg Leu	ttt Phe	ccg Pro	egg Arg 480	1440
30	ttg Leu	ggt Gly	gaa Glu	tcc Ser	gga Gly 485	gtc Val	act Thr	tct Ser	ggt Gly	ctc Leu 490	agc Ser	aaa Lys	ttg Leu	tac Tyr	tga 495		1485
35	<212	L> 49 2> Pl	T S	dopsi		halia	ana.										•
35	<213 <213 <213	L> 4! 2> Pl 3> Al	RT rabio	•	ls ti						• • • • • • • • • • • • • • • • • • • •	<i></i>	,	Tou	Cor	Leu	
	<213 <213 <213 <400 Met	L> 49 2> P1 3> A1 0> 6 Asp	RT rabio Leu	Gln	ls ti Arg	Thr	Leu			10					15		
40	<213 <213 <213 <400 Met 1 Leu	l> 49 2> P) 3> A 0> 6 Asp	RT rabio Leu Ile	Gln Lys 20	Arg 5 Thr	Thr	Leu Ala	Tyr	Asn 25	10 Glu	Lys	Gln	Leu	Phe 30	15 Gln	Pro	
	<211 <211 <211 <400 Met 1 Leu	L> 49 2> PP 3> As 0> 6 Asp Ile Glu	Thr	Gln Lys 20 Glu	Arg 5 Thr Asn	Thr Thr Ala	Leu Ala Asn	Tyr Ala 40	Asn 25 Met	10 Glu Thr	Lys Ala	Gln Val	Leu Met 45	Phe 30 Glu	15 Gln Arg	Pro Gly	
40	<211 <212 <213 <400 Met 1 Leu Leu Leu	L> 49 2> PR 3> AB 0> 6 ABP Lle Glu Lys 50	Leu Leu Ile Thr 35	Gln Lys 20 Glu Gln	Arg 5 Thr Asn Arg	Thr Thr Ala Arg	Leu Ala Asn Pro	Tyr Ala 40 Glu	Asn 25 Met His	10 Glu Thr Lys	Lys Ala Asn	Gln Val Ala 60	Leu Met 45 Tyr	Phe 30 Glu Ala	15 Gln Arg Thr	Pro Gly Met	
40	<211 <212 <213 <400 Met 1 Leu Leu Leu	L> 49 2> F) 3> A 0> 6 Asp Lle Glu Lys 50 Tyr	Leu Ile Thr 35 Thr	Gln Lys 20 Glu Gln Gly	Arg 5 Thr Asn Arg	Thr Thr Ala Arg Pro	Leu Ala Asn Pro 55 Arg	Tyr Ala 40 Glu Asp	Asn 25 Met His Tyr	10 Glu Thr Lys Glu	Lys Ala Asn Phe 75	Gln Val Ala 60 Tyr	Leu Met 45 Tyr Val	Phe 30 Glu Ala Ala	15 Gln Arg Thr	Pro Gly Met Arg 80	
40 45	<211 <212 <213 <400 Met Leu Leu Met 65 Val	L> 49 2> PR 3> As 0> 6 Asp Lle Glu Lys Tyr Leu	Leu Ile Thr 35 Thr Met	Gln Lys 20 Glu Gln Gly	Arg Thr Asn Arg Thr Ser	Thr Thr Ala Arg Pro 70 Leu	Leu Ala Asn Pro 55 Arg	Tyr Ala 40 Glu Asp	Asn 25 Met His Tyr	10 Glu Thr Lys Glu His 90	Lys Ala Asn Phe 75 Val	Gln Val Ala 60 Tyr Asp	Leu Met 45 Tyr Val Ala	Phe 30 Glu Ala Ala Asp	15 Gln Arg Thr Thr Ile 95	Pro Gly Met Arg 80 Val	
40 45 50	<211 <212 <213 <400 Met Leu Leu Met 65 Val	L> 49 2> PR 3> As 0> 6 Asp Lle Glu Lys 50 Tyr Leu Ile	Leu Ile Thr 35 Thr Met Ile	Gln Lys 20 Glu Gln Gly Arg	Arg 5 Thr Asn Arg Thr Ser 55	Thr Thr Ala Arg Pro 70 Leu Asp	Leu Ala Asn Pro 55 Arg Lys Val	Tyr Ala 40 Glu Asp Ser	Asn 25 Met His Tyr Leu Ile 105	Thr Lys Glu His 90 Asn	Lys Ala Asn Phe 75 Val	Gln Val Ala 60 Tyr Asp	Leu Met 45 Tyr Val Ala His	Phe 30 Glu Ala Ala Asp Ala 110	15 Gln Arg Thr Thr Ile 95 Leu	Pro Gly Met Arg 80 Val Glu	
40 45	<211 <211 <400 Met leu Leu Leu Met 65 Val Val	L> 49 2> PR 3> As 0> 6 Asp Ile Glu Lys 50 Tyr Leu Ile Glu	Leu Ile Thr 35 Thr Met Ile Ala Asp	Gln Lys 20 Glu Gln Gly Arg Ser 100 Gly	Arg 5 Thr Asn Arg Thr Ser Leu	Thr Thr Ala Arg Pro 70 Leu Asp	Leu Ala Asn Pro SS Aig Lys Val	Tyr Ala 40 Glu Asp Ser Pro Val 120	Asn 25 Met His Tyr Leu Ile 105 Arg	Thr Lys Glu His 90 Asn	Lys Ala Asn Phe 75 Val Trp Glu	Gln Val Ala 60 Tyr Asp Ile Asn	Leu Met 45 Tyr Val Ala His Leu 125	Phe 30 Glu Ala Ala Asp Ala 110 Glu	Arg Thr Thr Ile 95 Leu Asn	Pro Gly Met Arg 80 Val Glu Pro	
40 45 50	<211 <212 <400 Met 1 Leu Leu Met 65 Val Val Glu	L> 49 2> PR 3> As 0> 6 Asp Ile Glu Lys 50 Tyr Leu Glu Lys	Leu Ile Thr 35 Thr Met Ile Ala Asp 115	Gln Lys 20 Glu Gln Gly Arg Ser 100 Gly	Arg Thr Asn Arg Thr Ser B5 Leu Ala	Thr Thr Ala Arg Pro 70 Leu Asp Lys	Leu Ala Asn Pro SS Arg Lys Val Val	Tyr Ala 40 Glu Asp Ser Pro Val 120 Asp	Asn 25 Met His Tyr Leu Ile 105 Arg	Thr Lys Glu His 90 Asn Val	Lys Ala Asn Phe 75 Val Trp Glu Phe	Gln Val Ala 60 Tyr Asp Ile Asn Lys 140	Met 45 Tyr Val Ala His Leu 125 Leu	Phe 30 Glu Ala Ala Asp Ala 110 Glu Ser	Arg Thr Thr Ile 95 Leu Asn	Pro Gly Met Arg 80 Val Glu Pro Asn	
40 45 50	<211 <211 <400 Met 1 Leu Leu Leu Met 65 Val Val Glu Tyr	L> 49 2> PP 3> A 0> 6 Asp Ile Glu Lys 50 Tyr Leu Ile Glu Lys Leu Lys	Leu Ile Thr 35 Thr Met Ile Ala Asp 115 Lys	Gln Lys 20 Glu Gln Gly Arg Ser 100 Gly Gln	Arg Thr Asn Arg Thr Ser B5 Leu Ala Thr	Thr Thr Ala Arg Pro 70 Leu Asp Lys Asn Sex	Leu Ala Asn Pro SS Arg Lys Val Val Phe 135 Leu	Tyr Ala 40 Glu Asp Ser Pro Val 120 Asp	Asn 25 Met His Tyr Leu Ile 105 Arg Asn	Thr Lys Glu His 90 Asn Val Arg	Lys Ala Asn Phe 75 Val Trp Glu Phe Asp	Gln Val Ala 60 Tyr Asp Ile Asn Lys 140 Arg	Met. 45 Tyr Val Ala His Leu 125 Leu Val	Phe 30 Glu Ala Ala Asp Ala 110 Glu Ser Val	Arg Thr Thr Ile 95 Leu Asn Leu Met	Pro Gly Met Arg 80 Val Glu Pro Asn Leu 160 Cys	

```
Gly Gln Phe Cys Ala Val Phe Ile Asn Pro Cys Ile Phe His Thr Gly
                                        185
        Leu Phe Val Leu Gln Pro Ser Met Glu Val Phe Arg Asp Met Leu His
                                    200
   5
        Glu Leu Glu Val Lys Arg Asp Asn Pro Asp Gly Ala Asp Gln Gly Phe
                                215
       Leu Val Ser Tyr Phe Ser Asp Leu Leu Asn Gln Pro Leu Phe Arg Pro
                            230
       Pro Pro Asp Asn Arg Thr Ala Leu Lys Gly His Phe Arg Leu Pro Leu
                                                235
  10
                                            250
       Gly Tyr Gln Met Asp Ala Ser Tyr Tyr Tyr Leu Lys Leu Arg Trp Asn
                                        265
       Val Pro Cys Gly Pro Asn Ser Val Ile Thr Phe Pro Gly Ala Val Trp
                                    280
  15
       Leu Lys Pro Trp Tyr Trp Trp Ser Trp Pro Val Leu Pro Leu Gly Leu
                               295
       Ser Trp His His Gln Arg Arg Tyr Thr Ile Ser Tyr Ser Ala Glu Met
                           310
       Pro Trp Val Leu Thr Gln Ala Val Phe Tyr Leu Gly Ile Ile Leu Val
                                               315
 20
                                           330
       Thr Arg Leu Ala Arg Pro Asn Met Thr Lys Leu Cys Tyr Arg Arg Ser
                                       345
      Asp Lys Asn Leu Ser Met Ile Gln Thr Ala Phe Lys Phe Val Ala Leu
                                   360
      Leu Phe Ile Leu Ser Ala Tyr Ile Ile Pro Phe Phe Ile Ile Pro Glr
 25
                               375
      Thr Ile His Pro Leu Ile Gly Trp Ser Leu Tyr Leu Thr Gly Ser Phe
                                                   380
                          390
                                               395
      Ala Leu Ser Thr Ile Pro Ile Asn Ala Phe Leu Leu Pro Ile Leú Pro
 30
                      405
                                          410
      Val Ile Thr Pro Trp Leu Gly Ile Phe Gly Thr Leu Leu Val Met Ala
                  420
                                      425
      Phe Pro Ser Tyr Pro Asp Gly Val Val Arg Ala Leu Ser Val Phe Gly
                                - 440
      Tyr Ala Phe Cys Cys Ala Pro Phe Leu Trp Val Ser Phe Val Lys Ile
                                                      445
                              455
                                                  460
      Thr Ser His Leu Gln Ile Met Ile Asp Lys Glu Val Leu Phe Pro Arg
                          470
     Leu Gly Glu Ser Gly Val Thr Ser Gly Leu Ser Lys Leu Tyr
                                              475
                                                                  480
40
                      485
                                          490
     <210> 7
45
     <211> 1494
     <212> DNA
     <213> Arabidopsis thaliana
     <220>
50
     <221> CDS
     <222> (1)..(1494)
     atg gat ttg cag aga ggc ttt gtg ttc ttg tct ttg gtt cta tct ttt
     Met Asp Leu Gln Arg Gly Phe Val Phe Leu Ser Leu Val Leu Ser Phe
55
    atg ata atc gaa acg acg gcg tat cga gag aga cag ctg ctg ctg
    Met Ile Ile Glu Thr Thr Ala Tyr Arg Glu Arg Gln Leu Leu Leu
60
                                      25.
    caa cca ccg caa gaa acg gcg ata gat acc gca aac gcg gtg gtg acg
```

		_							1-4		•				
			- •				¥0					45		al Thr	
5		50	•			55		ALG	Arg	KIO (60 GTØ H	is L	ys A	ac gca sn Ala	
10	65				70	1400	GLY	IHE	PIQ	75	Asp T	Ài G	Lu Pi	c tac e Tyr . 80	240
15				85		A-4-C ,	ara	ser	90 Leu	Arg S	er L	eu Hi	5V a. e	g gaa 1 Glu 5	288
	gct g Ala A		100		***		Jer .	105	авр (Val P	ro Le	u Ar 11	g Tr O	p Val	336
20	caa ad Gln Ti	11.	5	4-4		j Jej	120	MIG .	rže /	al V	al Ar 12	g Va 5	l Gl	u Asn	384
25		0		,***	1	.35	7111 7	var s	asn P	ne Ar 14	sn Se 10	r Ar	y Phe	Lys	432
30	ctt ac Leu Th 145		- ******	Lys :	150	Ar w	na 1	xp A	ua L 1	eu Se 55	er Asj	, EKL d	Ası	Arg. 160	480
35	gtg gt Val Va		. 204	165	nia n	вр и	sn L	eu p 1	ne L 70	eu Ly	s Ļķ	3 Ala	175	Glu	528
	ttg tt		180	GIY A	ng r	ne C	ys A	1a V 85	al Pi	ie Il	e Asr	190	Cys	Ile	576
40	ttc cae Phe His	e act Thr 195	ggt Gly	ctc t Leu F	tc gr	at he	tg ca eu Gi	ag c	ca to ro Se	a gt er Va	g gaa 1 Glu 205	Val	ttc Phe	aag Lys	624
45	gac ato Asp Met 210		cat (gag c Glu L	ta ca eu Gl 21	TIT AS	t gg	ga aq ly A	ga aa	ig aai 8 Asi 22	r Pro	gat Asp	gga Gly	gct Ala	672
50	gat caa Asp Glr 225	ggt Gly	ttt (Phe 1	ren A	tc aç al Se 30	gt ta er Ty	c tt r Ph	c to le Se	et ga er As 23	b res	ctt Leu	gac Asp	caa Glņ	cct Pro 240	720
55	ctc ttt Leu Phe	agt Ser	ATO 1	ecg a Pro s 145	gt aa er As	ic gg	a to y Se	t gt r Va 25	il Le	t aat u Asr	ggt Gly	cac His	ttg Leu 255	aga Arg	768
	ett eee Leu Pro	tta Leu	ggc t Gly 1 260	ac c	aa at In Me	g ga t As	c gc p Al 26	a Se	t ta r Ty	t tto r Phe	tat Tyr	ctt Leu 270	aag Lys	cta Leu	816
60	aga tgg Arg Trp	aac Asn 275	ata c Ile P	ce to	gt gg Ys Gl	a cc y Pr 28	o As	c ag	t gt r Va	g att l Ile	aca Thr 285	ttc Phe	eeg Pro	gga Gly	864
				•											

	get gtt tgg tta aag cca tgg tac tgg tgg tca tgg cct gtt ctt cca Ala Val Trp Leu Lys Pro Trp Tyr Trp Trp Ser Trp Pro Val Leu Pro 290 295 300	•
· 4	cta ggt ttc tca tgg cac gag cag cgt cgc gcc act ata ggg tac tca Leu Gly Phe Ser Trp His Glu Gln Arg Arg Ala Thr Ile Gly Tyr Ser 305 310 315 320	960
•	Ala Glu Met Pro Leu Val Ile. Ile Gln Ala Met Phe Tyr Leu Gly Ile 325	1008
1:	340 See Ala Arg Pro Asn Ile Thr Lys Leu Cys Tyr	1056
20	360 365 Leu	1104
25		1152
	atc cct cac act atc cac cca ctc atc ggc tgg tcg ctc tac ttg atg Ile Pro His Thr Ile His Pro Leu Ile Gly Trp Ser Leu Tyr Leu Met 385 390 395 400	1200
30	Ala Ser Phe Ala Leu Ser Ser Ile Ser Ile Asn Thr Leu Leu Pro 405 410	1248
35	acg ctc cct gtt ctc act cca tgg cta gga att ctc ggc act ctc ctt Thr Leu Pro Val Leu Thr Pro Trp Leu Gly Ile Leu Gly Thr Leu Leu 420 425 430	1296
40	gtc atg gcc ttc cct tgg tac cct gat gga gtg gtc aga gcc ttg tca Val Met Ala Phe Pro Trp Tyr Pro Asp Gly Val Val Arg Ala Leu Ser 435	1344
45	450 455 460	1392
	470 475 Glu Val Leu	1440
50	ttc ccg cga ttg gga gac tca ggg gtc act tca ggc ttg	L 488
55	tat tag	494
60	<210> 8 <211> 497 <212> PRT <213> Arabidopsis thaliana	

	76	
	<400> 8 , Met Asp Leu Gla Are Gla Plane	
	Met Asp Leu Gln Arg Gly Phe Val Phe Leu Ser Leu Val Leu Ser Ph	ie
	5 20 Leu	73
	Gln Pro Pro Gln Glu Thr Ala Ile Asp Thr Ala Asn Ala Val Val Th	_
	Val Gln Asp Arg Gly Lev Term The 45	r
10	Val Gln Asp Arg Gly Leu Lys Thr Arg Arg Pro Glu His Lys Asn Al	a
,,	65 Tyr Met Gly Thr Pro Arg Asp Tyr Glu Phe Tyr	E:
	Val Ala Thr Arg Val Leu Ile Arg Ser Leu Arg Ser Leu His Val Gly)
, .	Ala Asp Leu Val Val Tia Nia - 95	1
15	100 105 110	L
	Gin Thr Leu Glu Glu Glu Asp Gly Ala Lys Val Val Arg Val Glu Asn	Ł
	Val Asp Asn Pro Tyr Arg Arg Gln Thr Asn Phe Asn Ser Arg Phe Year	
20	Leu Thr Leu Asn Lys Leu Tyr Ala Trp Ala Leu Ser Asp Tyr Asp Arg	•
•	Val Val Mor Lou Age 210 155 156	
	Val Val Met Leu Asp Ala Asp Asn Leu Phe Leu Lys Lys Ala Asp Glu	
25	then Pole Gin Cys Gly Arg Phe Cys Ala Val Phe Ile Asn Pro Cys Ile	
	Phe His Thr Gly Leu Phe Val Leu Gln Pro Ser Val Glu Val Phe Lys	
	Asp Met Leu His Glu Leu Glu Vol Glu Vol Glu Vol	
30	Asp Met Leu His Glu Leu Gln Val Gly Arg Lys Asn Pro Asp Gly Ala 210 215 220 220	
	Asp Gln Gly Phe Leu Val Ser Tyr Phe Ser Asp Leu Leu Asp Gln Pro	
	Leu Phe Ser Pro Pro Ser Asn Gly Ser Val Leu Asn Gly His Leu Arg	
35	Leu Pro Leu Gly Tyr Gln Met Asp Ala Ser Tyr Phe Tyr Leu	
33	Arg Tro Asn Tle Pro Cue Cla Date 265 270	
	Arg Trp Asn Ile Pro Cys Gly Pro Asn Ser Val Ile Thr Phe Pro Gly 275 280 285	
	Ala Val Trp Leu Lys Pro Trp Tyr Trp Trp Ser Trp Pro Val Leu Pro 290 295 300	
40	Leu Gly Phe Ser Trp His Glu Gln Arg Arg Ala Thr Ile Gly Tyr Sor	
	Ala Glu Met Pro Leu Val Ile Ile Gln Ala Met Phe Tyr Leu Gly Ile	
	Ile Val Val The last to 330 335	
45	Ile Val Val Thr Arg Leu Ala Arg Pro Asn Ile Thr Lys Leu Cys Tyr	
	Arg Arg Ser Asp Arg Asp Leu Thr Thr Ile Gln Ala Gly Phe Lys Leu	
•	Ile Ala Leu Leu Ser Val Val Ala Ala Tur Ile Phe Pro Pho Pro	
50	370 375 Ile Pro His Thr Ile His Pro Leu Ile Gly Trp Ser Leu Tyr Leu Met 385 390	
	390 395 Leu Tyr Leu Met	
	Ala Ser Phe Ala Leu Ser Ser Ile Ser Ile Asn Thr Leu Leu Pro 405 410	
55	Thr Leu Pro Val Leu Thr Pro Tro Leu Gly Ile Leu Gly Thr Leu Leu	
	Val Met Ala Phe Pro Trp Tyr Pro Asp Gly Val Val Arg Ala Leu Ser	
•		
60	Val Phe Ala Tyr Ala Phe Cys Cys Ala Pro Phe Val Trp Val Ser Phe 450 455 460	
60	Arg Lys lie Thr Ser His Leu Gln Val Leu Ile Glu Lys Glu Val Leu	
	Phe Pro Arg Leu Gly Asp Ser Gly Val Thr Ser Gly Phe Ser Lys Leu	

	. 1	ľyr			4	185				4	90		•		4	95		
;	5																	
•		210:	· 9 · 194															
			DNA															
	<				psis	tha	lian	a										
10)				~			_										
		220>																
			CDS			•												
	ج,	422>	(1)	(1	944)													
15	<	400>	9															
	at	ta c	at c	ca q	et to	ca to	rt ac	rt ~+	٠~ +-	·n _L						t aat		
	Me	et H	is P	ro Al	La Se	57. CA	78 Se	r Le	n Se	a ct	ic ac	gg aa	aa gt	c aa	a ct	t aat u Asr	48	
		1				5				1	0	g L	Aa As	of PA	s Le	u Asr	l	
20															1			
20	TI	.C CI	בנ פני מוז איני	a gr	g ag	a at	g aa	g ct	c tc	t tc	t ga	a aç	jt ac	c at	g ge	g ccg	96	•
		116	314 HI		O TE	g me	r ry	s Le	u se	r se	r Gl	u Se	r Pi	o Me	E Ala	g ccg	,	
				4	. •				2	5				3		-		
~=	to	a ag	rt ca	g to	a ag	t ca	t ca	a ct	t ta	c at	t t~	c 20	+			a aaa		
25	Se	x Se		-	r Se	r Hi	s Ar	g Le	u Ty	r Il	e Se	r Se	r Gi	g aas	a aca	a aaa : Lys	144.	
			3	5				4	0				4		, 1111	. пАв	•	
	20	~ ~~	~ ~~			_												
	Th	r Lv	s Ar	a LL	e ca	a aga	a aad	998	a tac	act	t ct	c ga	t gt	t gaa	atg	r ,tgt	192	
30		, 5	0	3	C GI	u AL	J ASI 55	رحب م	Ty	r Thi	c Le	u As	p Va	t gaa l Glu	Met	Сув		
												6	_					
	gt	aa	c tt	c te	t agi	t ctg	aaa	ctt	gtt	tto	, tti	. ct:	t ato	g atg	ata		•••	
	Va.	L As:	n Ph	e Sei	r Sei	r Let	Lys	Leu	Val	. Leu	Phe	Le	ı Met	g atg : Met	Ten	ycc Val	240	
35	65	>				70)				75	;				80		
	act	. ato	7 tt/	. 202	050													
	Ala	Met	Phe	The	Lev	. cac	· Cve	Cor	CCA	ccg	ttg	caa	att	cct Pro	gaa	gat	288	
					85	x-	~	Ger	FIQ	90 90	nen	GLI	1 Ile	Pro		Asp		
40															95			
40	CCa	tca	agt	ttt	gca	aac	aaa	tgg	ata	cta	gaa	cct	get	gta	acc	aca	336	
	Pro	sei	Ser			Asn	Lys	Trp	TTE	Leu	Glu	Pro	Ala	gta Val	Thr	Thr	330	
				100	,				105					110				
	gat	cct	ago	tat	ata	get	aca	ter	~~	-t-				agt.				
45	Āsp	Pro	Arg	Tyr	Ile	Ala	Thr	Ser	Glu	Ile	Aan	regg	ABC	agt Ser	atg	tca	384	
			115					120			41044		125	ser	wet	ser		
	a++																	
	TAN	yr. Val	yrt Val	gag	Cat	tac	tta	tct	aac	aga	agc	gag	tat	caa	gga	att	432	
50	204	130		GIU	uta	TYT	135	seŗ	Gly	Arg	Ser	Glu	Tyr	caa Gln	Gly	Ile		
												140						
	ggc	ttt	cta	aat	ctc	aac	gat	aac	gag	att	aat	CGO	taa	cag				
		Phe	Leu	Asn	Leu	Asn	asp	Asn	Glu	Ile	Asn	Arg	Tro	cag Gln	ycy Val	gcc Tat	480	
55	145				•	150					155					160		
33	ate	200	+															
	Tle	T.va	Sar	uic	rgr Cva	cag	cat	ata	gct	ttg	cat	cta	gac	cat	gct	gca	528	
		~J =	JET.	****	165	GTI	n15	тте	ALA	ren	His	Leu	Asp	His :	Ala .	Ala	•	
•							•			170					175			
60	agt	aa¢	ata	act	tgg	aaa	tct	tta	tac	cca	ga2	taa	att	gac (~~	~~~	E76	
	ser	Asn	Ile		Trp	Lys	Ser :	Leu	Tyr	Pro	Glu	Tro.	:Ile	gac (Asp (344 ! 31:	944 514 -	576	
				180					185			-		190				
														•				

	gaa aaa ttc aaa gtc ccc act tgt cct tct ctt cct tgg att caa gtt 624 Glu Lys Phe Lys Val Pro Thr Cys Pro Ser Leu Pro Trp Ile Gln Val 200 205	ŧ
	cct gac aag tct cga atc gat ctt atc att gcc aag ctc cca tgt aac 672 Pro Asp Lys Ser Arg Ile Asp Leu Ile Ile Ala Lys Leu Pro Cys Asn 210 215	ı
10	Lys Ser Gly Lys Trp Ser Arg Asp Val Ala Arg Leu His Leu Gln Leu 230 235 240	
15	245 250 255	
20	265 270	
25		
•	ctt cac cag tta aga caa aag tta caa ctt cct gtt ggt tcc tgt gaa 912 Leu His Gln Leu Arg Gln Lys Leu Gln Leu Pro Val Gly Ser Cys Glu 290 295 300	
30	ctt tct gtt cct ctt caa gct aaa gat aat ttc tac tcg gca aat gcc 960 Leu Ser Val Pro Leu Gln Ala Lys Asp Asn Phe Tyr Ser Ala Asn Ala 310 315 320	
35	aag aaa gaa gog tac gog acg atc ttg cac tca gat gat gct ttt gtc 1008 Lys Lys Glu Ala Tyr Ala Thr Ile Leu His Ser Asp Asp Ala Phe Val 325 330 335	
40	tgt gga gcc att gca gta gca cag agc att cga atg tca ggc tct act 1056 Cys Gly Ala Ile Ala Val Ala Gln Ser Ile Arg Met Ser Gly Ser Thr 340 345 350	
45	cgc aat ttg gta ata cta gtc gat gat tcg atc agt gaa tac cat aga 1104 Arg Asn Leu Val Ile Leu Val Asp Asp Ser Ile Ser Glu Tyr Ris Arg 355 360 365	
	agt ggc ttg gaa tca gct gga tgg aag att cac aca ttt caa aga atc 1152 Ser Gly Leu Glu Ser Ala Gly Trp Lys Ile His Thr Phe Gln Arg Ile	
50	370 375 380	
ee	aga aac ccg aaa gct gaa gca aat gca tat aac caa tgg aac tac agc 1200 Arg Asn Pro Lys Ala Glu Ala Asn Ala Tyr Asn Gln Trp Asn Tyr Ser 385 390 395 400	
55	aaa ttc cgt ctt tgg gaa ttg aca gaa tac aac aag atc atc ttc att 1248 Lys Phe Arg Leu Trp Glu Leu Thr Glu Tyr Asn Lys Ile Ile Phe Ile 405 410 415	
60	gat gca gac atg ctt atc ctc aga aac atg gat ttc ctc ttc gag tac 1296 Asp Ala Asp Met Leu Ile Leu Arg Asn Met Asp Phe Leu Phe Glu Tyr 420 425 430	

	ccc gaa atc tcc aca act gga aac gac ggt acg ctc ttc aac tcc ggt Pro Glu Ile Ser Thr Thr Gly Asn Asp Gly Thr Leu Phe Asn Ser Gly 435	1344
	5 cta atg gtg att gaa cca tca aat tca aca ttc cag tta cta atg gat Leu Met Val Ile Glu Pro Ser Asn Ser Thr Phe Gln Leu Leu Met Asp 450 455 460	1392
11	465 Ash Asp 11e Ash Ser Tyr Ash Gly Gly Asp Gln Gly Tyr Leu 475 476	1440
15	495 490 495	1488
20	720	1536
25	acg aga cta ttt gga gct gat cct ccg ata ctc tac gtt ctt cat tac Thr Arg Leu Phe Gly Ala Asp Pro Pro Ile Leu Tyr Val Leu His Tyr 525	1584
20	cta ggt tac aac aaa cca tgg gta tgc ttc aga gac tac gat tgc aat Leu Gly Tyr Asn Lys Pro Trp Val Cys Phe Arg Asp Tyr Asp Cys Asn 530 535 540	1632
30	Tro Asn Val Val Gly Tyr His Gln Phe Ala Ser Asp Glu Ala His Lys 545 550 550 560	1680
35	act tgg tgg aga gtg cac gac gcg atg cct aag aaa ttg cag agg ttt Thr Trp Trp Arg Val His Asp Ala Met Pro Lys Lys Leu Gln Arg Phe 565 570 575	1728
40	tgt cta ctg agt tcg asa caa aag gcg caa ctt gag tgg gat cgg aga Cys Leu Leu Ser Ser Lys Gln Lys Ala Gln Leu Glu Trp Asp Arg Arg 580 585	1776
ΛE	595 . 600 GOS TIP ATG ILE LYS ILE	1824 ´
45	610 615 File Gitt Asp Phe Cys Phe Trp Glu	1872
50	625 . 630 635 640	1920
55	gct gat tct tcc tcc acc gcg taa Ala Asp Ser Ser Thr Ala 645	944
60	<210> 10 <211> 647 <212> PRT <213> Arabidopsis thaliana	

		0> 1														
	1.		Pro		5					10				_	15	
5			Thr	20					25					30		
			Gln 3:5					40					45			_
40		50	Arg				55					60				-
10	65		Phe			70					75				•	80
			Phe		85					90					95	-
15			Ser	100					105			•		110		
			Arg					120				_	125			
20		130	Val Leu				135		-			140	-		_	•
	145					150					155					160
			Ser		165					170			-		175	
25			Ile	180					185					190		
			Phe 195				•	200					205			
20		210	Lys				215					220			-	
30	225		Gly	_	_	230	_	_			235					240
			Ala		245					250	_		•	-	255	
35	•		Leu	260			•		265					270		_
			275 Gln			_		280			_		285	-		_
40		290	Val		-		295					300	-			
70	305		Glu			310					315					320
	-	-	Ala		325			•		330		_			335	
45	•	-		340					345		•			350		
			355 Leu					360					365			
50		370	Pro				375					380				
	385		Arg	-		390				_	395		_			400
•	_		Asp		405					410					415	
55	Pro	Glu	Ile	420 Ser	Thr	Thr	Gly	Asn	425 Asp	Gly	Thr	Leu	Phe	430 Asn	ser	Gly:
	Leu	Met	435 Val	Ile	Glu	Pro	Ser	440 Asn	ser	Thr	Phe	Gln	445 Leu	Leu	Met	Asp
60		450	Asn				455					460				Leu
	465 Asn	Glu	Ile	Phe	The	470 Trp	Trp	His	Arg	Ile	475 Pro	Lys	His	Met	Asn	480 Phe
	•															

		• • • • • • • • • • • • • • • • • • • •
	485 490 495 Leu Lys His Phe Trp Glu Gly Asp Thr Pro Lys His Arg Lys Ser Ly	
5	500 505 510 Thr Arg Leu Phe Gly Ala Asp Pro Pro Ile Leu Tyr Val Leu His Ty	T.
	515 520 525 Leu Gly Tyr Asn Lys Pro Trp Val Cys Phe Arg Asp Tyr Asp Cys As 530 535	n
·	Trp Asn Val Val Gly Tyr His Gln Phe Ala Ser Asp Glu Ala His Ly	3
10	Thr Trp Trp Arg Val His Asp Ala Met Pro Lys Lys Leu Gln Arg Ph	0 e
	Cys Leu Leu Ser Ser Lys Gln Lys Ala Gln Leu Glu Trp Asp Arg Ar	5
15	Gln Ala Glu Lys Ala Asn Tyr Arg Asp Gly His Trp Arg Ile Lys Il 595 600 605	e
	Lys Asp Lys Arg Leu Thr Thr Cys Phe Glu Asp Phe Cys Phe Trp Gl 610 620	n
	Ser Met Leu Trp His Trp Gly Glu Thr Gln Thr Asn Ser Thr Val Al.	
20	Ala Asp Ser Ser Ser Thr Ala 645	J
25	<210> 11	
	<211> 1857 <212> DNA <213> Arabidopsis thaliana	
30	<220>	
	<221> CDS <222> (1)(1857)	
	<400> 11	
35	atg ata cet tee tea agt eee atg gag tea aga eat ega ete teg tee Met Ile Pro Ser Ser Pro Met Glu Ser Arg His Arg Leu Ser Phe	48
	1 5 10 15	
40	tca aaa gag aag aca agt agg agg aga ttt caa aga att gag aag ggt Ser Lys Glu Lys Thr Ser Arg Arg Phe Gln Arg Ile Glu Lys Gly	96
	20 . 25 30	
45	gtc aag ttc aac act ctg aaa ctt gtg ttg att tgt ata atg ctt gga Val Lys Phe Asn Thr Leu Lys Leu Val Leu Ile Cys Ile Met Leu Gly 35 40	144
	got tig the acg ate tae egt the egt tat eea eeg eta caa att eet	. 192
	Ala Leu Phe Thr Ile Tyr Arg Phe Arg Tyr Pro Pro Leu Gln Ile Pro	,
50	gaa att cca act agt tit ggt cit act act gat cct cgc tat gta got	240
	Glu Ile Pro Thr Ser Phe Gly Leu Thr Thr Asp Pro Arg Tyr Val Ala 65 70 75 80	
55	aca get gag ate aac tgg aac cat atg tea aat ett gtt gag aag cae	288
	Thr Ala Glu Ile Asn Trp Asn His Met Ser Asn Leu Val Glu Lys His 85 90 95	
60	gta ttt ggt aga agc gag tat caa gga att ggt ctt ata aat ctt aac	
09,	Val Phe Gly Arg Ser Glu Tyr Gln Gly Ile Gly Leu Ile Asn Leu Asn 100 105 110	

	gat Asp	aac Asn	gag Glu 115	att Ile	gat Asp	cga Arg	ttc Phe	aag Lys 120	Glu	gta Val	acg Thr	rys aaə	tet Ser 125	gac	tgt Cys	gat Asp	384 _.
5	cat His	gta Val 130	gct	ttg Leu	cat His	cta Leu	gat Asp 135	Tyr	gct Ala	gca Ala	aag Lys	aac Asn 140	ata Ile	aca Thr	tgg T r p	gaa Glu	432
10	tet Ser 145	tta Leu	tac Tyr	ecg Pro	gaa Glu	tgg Trp 150	att Ile	gat Asp	gaa Glu	gtt Val	gaa Glu 155	gaa Glu	ttc Phe	gaa Glu	gtc Val	cct Pro 160.	480
15	act Thr	tgt Cys	ect Pro	tct Ser	ctg Leu 165	cct Pro	ttg Leu	att Ile	Caa Gln	att Ile 170	cct Pro	ggc Gly	aag Lys	cct Pro	cgg Arg 175	att Ile	528
20	gat Asp	ctt Leu	gta Val	att Ile 180	gcc Ala	aag Lys	ctt Leu	ccg	tgt Cys 185	gat Asp	aaa Lys	tca Ser	Gly	aaa Lys 190	tgg Trp	tct Ser	576
	aga Arg	gat Asp	gtg Val 195	gct Ala	Arg	ttg Le u	cat His	tta Leu 200	caa Gln	ctt Leu	gca Ala	gca Ala	gct Ala 205	cga Arg	gtg Val	gcg Ala	624
25	get Ala	tct Ser 210	tct Ser	aaa Lys	gga Gly	ctt Leu	cat His 215	aat Asn	gtt Val	cat His	Val	att Ile 220	ttg Leu	gta Val	tet Se r	gat Asp	672
30	Cys 225	ttt Phe	cca Pro	ata Ile	Pro Pro	aat Asn 230	ctt Leu	ttt Phe	acg Thr	gjy ggt	caa Gln 235	gaa Glu	ctt Leu	gtt Val	gcc Ala	cgt Arg 240	720
35	caa Gln	ely gga	aac Asn	Ile	tgg Trp 245	ctg Leu	tat Tyr	aag Lys	cct Pro	aat Asn 250	ctt Leu	cac His	Glņ Cag	cťa Leu	aga Arg 255	caa Gln	· 768
40	Lys	Leu	Gln	Leu 260	Pro	gtt Val	Gly	ser	Сув 265	Glu	Leu	Ser	Val	Pro 270	Leu	Gln	816
45	gct Ala	Lys	gat Asp 275	aat Asn	ttc Phe	tac Tyr	ser .	gca Ala 280	ggt Gly	gca Ala	aag Lys	Lys	gaa Glu 285	gct Ala	tac Tyr	gcg Ala	864
•	act Thr	atc Ile 290	ttg Leu :	cat His	tct Ser	Ala	caa Gln 295	ttt Phe	tat Tyr	gtc Val	Cys	gga Gly 300	gcc	att Ile	gca Ala	gct Ala	912
50	gca Ala 305				Arg .					Thr					İle		960
	gtt Val			Thr					His								1008
60	gga Gly		Lys			atg : Met :		Gln					Pro				1056
	cca	aat	gc¢	tac	aac	gaa	tg g	aac	tac	agç	aag	ttt	cgt	ctt	tgg	caa	1104

	23	
	Pro Asn Ala Tyr Asn Glu Trp Asn Tyr Ser Lys Phe Arg Leu Trp Gln 355 360 365	~-
	ctg act gaa tac agt aag atc atc ttc atc gat gca gac atg ctt atc 5 Leu Thr Glu Tyr Ser Lys Ile Ile Phe Ile Asp Ala Asp Met Leu Ile 370 375 380	1152
10	395 400	1200
15	•=•	1248
	tct aat tca aca ttc cag tta cta atg gat aac att aat gaa gtt gtg Ser Asn Ser Thr Phe Gln Leu Leu Met Asp Asn Ile Asn Glu Val Val 420 425 430	1296
20	tot tac aac gga gga gac caa ggt tac ott aac gag ata tto aca tgg Ser Tyr Asn Gly Gly Asp Gln Gly Tyr Leu Asn Glu Ile Phe Thr Trp 435 440	1344
25	tgg cat cgg att cca aaa cac atg aat ttc ttg aag cat ttc tgg gaa Trp His Arg Ile Pro Lys His Met Asn Phe Leu Lys His Phe Trp Glu 450 455 460	1392
30	465 470 475 Leu Phe Gly Ala	1440
35	485 490 495	1488
	500 505 Trp Asn Val Asp Ile Phe	1536
40	cag gaa ttt gct agt gac gag gct cat aaa acc tgg tgg aga gtg cac 1 Gln Glu Phe Ala Ser Asp Glu Ala His Lys Thr Trp Trp Arg Val His 515 520 525	.584
45	gac gca atg cct gaæ aac ttg cat aag ttc tgt cta cta aga tcg aaa 1 Asp Ala Met Pro Glu Asn Leu His Lys Phe Cys Leu Leu Arg Ser Lys 530 535	632
50	545 550 555 560	680
55	565 570 575	728
	act tgt ttc gaa gat ttc tgc ttt tgg gag agt atg ctt tgg cat tgg Thr Cys Phe Glu Asp Phe Cys Phe Trp Glu Ser Met Leu Trp His Trp 580 585 590	776
60		324
	Gly Glu Thr Asn Ser Thr Asn Asn Ser Ser Thr Thr Thr Thr Ser Ser	. =

ccg ccg cat aaa acc gct ctc cct tcc ctg tga Pro Pro His Lys Thr Ala Leu Pro Ser Leu <210> 12 <211> 618 <212> PRT <213> Arabidopsis thaliana Met Ile Pro Ser Ser Ser Pro Met Glu Ser Arg His Arg Leu Ser Phe Ser Lys Glu Lys Thr Ser Arg Arg Arg Phe Gln Arg Ile Glu Lys Gly Val Lys Phe Asn Thr Leu Lys Leu Val Leu Ile Cys Ile Met Leu Gly Ala Leu Phe Thr Ile Tyr Arg Phe Arg Tyr Pro Pro Leu Gln Ile Pro Glu Ile Pro Thr Ser Phe Gly Leu Thr Thr Asp Pro Arg Tyr Val Ala Thr Ala Glu Ile Asn Trp Asn His Met Ser Asn Leu Val Glu Lys His Val Phe Gly Arg Ser Glu Tyr Gln Gly Ile Gly Leu Ile Asn Leu Asn Asp Asn Glu Ile Asp Arg Phe Lys Glu Val Thr Lys Ser Asp Cys Asp His Val Ala Leu His Leu Asp Tyr Ala Ala Lys Asn Ile Thr Trp Glu Ser Leu Tyr Pro Glu Trp Ile Asp Glu Val Glu Phe Glu Val Pro Thr Cys Pro Ser Leu Pro Leu Ile Gln Ile Pro Gly Lys Pro Arg Ile Asp Leu Val Ile Ala Lys Leu Pro Cys Asp Lys Ser Gly Lys Trp Ser Arg Asp Val Ala Arg Leu His Leu Glo Leu Ala Ala Ala Arg Val Ala Ala Ser Ser Lys Gly Leu His Asn Val His Val Ile Leu Val Ser Asp Cys Phe Pro Ile Pro Asn Leu Phe Thr Gly Gln Glu Leu Val Ala Arg Gln Gly Asn Ile Trp Leu Tyr Lys Pro Asn Leu His Gln Leu Arg Gln Lys Leu Gln Leu Fro Val Gly Ser Cys Glu Leu Ser Val Pro Leu Gln Ala Lys Asp Asn Phe Tyr Ser Ala Gly Ala Lys Lys Glu Ala Tyr Ala Thr Ile Leu His Ser Ala Gln Phe Tyr Val Cys Gly Ala Ile Ala Ala Ala Gln Ser Ile Arg Met Ser Gly Ser Thr Arg Asp Leu Val Ile Leu Val Asp Glu Thr Ile Ser Glu Tyr His Lys Ser Gly Leu Val Ala Ala Gly Trp Lys Ile Gln Met Phe Gln Arg Ile Arg Asn Pro Asn Ala Val Pro Asn Ala Tyr Asn Glu Trp Asn Tyr Ser Lys Phe Arg Leu Trp Gln Leu Thr Glu Tyr Ser Lys Ile Ile Phe Ile Asp Ala Asp Met Leu Ile

										23							
	Leu 385	Arg	Asn	Ile	Asp	Phe 390		Phe	Glu	Phe	Pro		Ile	Ser	Ala	Thr 400	
	Gly	Asn	Asn	Ala	Thr 405			Asn	. ser	Gly 410	Leu		Val	Val	Glu 415	Pro	
5	Ser	Asn	Ser	Thr 420	Phe	Gln	Leu	Leu	Met 425	Asp		Ile	Asn	Glu 430	Val	Val	
	Ser	Tyr	Asn 435	Gly	Gly	Asp	Gln	Gly 440			Asn	Glu	Ile 445	Phe	Thr	Trp	
10	Trp	His 450	Arg	Ile	Pro	гàа	His 455		Asn	Phe	Leu	Lys 460		Phe	Trp	Glu	
	Gly 465	Asp	Glu	Pro	Glu	Ile 470	Lys	Lys	Met	Lys	Thr 475		Leu	Phe	Gly	Ala 480	
					Leu 485					490	Leu				495	Pro	
15				500	Arg				505					510	Ile		
			515		Ser			520					525				
20		530			Glu		535					540		•			
	545				Leu	550					555			_	-	560	
					His 565					570				•	ミウに	Lys	
25				200					282					590		Trp ,	
			595		Ser			600			Thr	Thr	Thr 605	Thr	Ser	Ser	
30	Pro	Pro 610	His	Lys	Thr	Ala	Leu 615	Pro	Ser	Leu				. ,	•		
													•				
	<210)> 13	;										•				
35	<21]	> 21 > DN	.30														
	<213	> Az	abid	tagol	s th	alia	na										
40	<220 <221)> L> CID	s														
	<222	!> (1	.) (1980))												
		> 13		tet	ccc	act	act	cct	GC3	·		200	202	200	aat-	aat-	48
45	Met	Ala	Asn	Ser	2x0 5	Ala	Ala	Pro	Ala	Pro 10	Thr	Thr	Thr	Thr	Gly 15	Gly	10
		taa	000	~~		ata	- ee	6 50	•			5 00				D. 65	96
50					Arg												76
0.0	aga	tte	caa		aat	age	aaa	gga		aac	acta	tea	gat.	•	at.a	222	144
	Arg																
55	cca	+++		atr	ata	aat'	+++		202	caa	asc	222		ant	agt	tat	192
	_				Ile			_			_			_	_	_	270
60	+~+		++~	200	aag	ttt	•	at-c	ata	227	ct+	•	tta	+++	at.c	att	240
······································					Lys												- ·
	03					,,,					10					4 0	

5	ctc Leu	tct Ser	gcc Ala	act Thr	ctc Leu 85	ttc Phe	acc Thr	att Ile	atç Ile	tat Tyr 90	tct Ser	cct Pro	gaa Glu	gct Ala	tat Tyr 95	cat His	288
	cat His	tct Ser	ctt Leu	tcc Ser 100	cac His	tca Ser	tct Se r	tct Ser	cga Arg 105	tgg Trp	ata Ile	tgg Trp	aga Arg	aga Arg 110	caa Gln	gat Asp	336
10	cca Pro	cgt Arg	tac Tyr 115	ttc Phe	tcg Ser	gat Asp	ctg Leu	gat Asp 120	ata Ile	aac Asn	tgg Trp	gac Asp	gat Asp 125	gtg Val	act Thr	aaa Lys	384
15	acc Thr	ctt Leu 130	gag Glu	aac Asn	atc Ile	gaa Glu	gaa Glu 135	eja aac	cgt Arg	acg Thr	atc Ile	ggt Gly 140	gtc Val	ttg Leu	aat Asn	ttt Phe	432
20	gat Asp 145	tcg Ser	aac Asn	gag Glu	atc Ile	caa Gln 150	cga Arg	tgg Trp	aga Arg	gaa Glu	gta Val 155	tcc Ser	aag Lys	agc Ser	aag Lys	gac Asp 160	480
25						aaa Lys											528
						gac Asp									Ģlu		576
30	caa Gln																624
35						gat Asp		Ile									672
40	gaa Glu 225																720
45	gct Ala																768
						tgc Cys											816
50	gat Asp					aga Arg											864
55						aag Lys											912
60	tct Ser 305	ctt Leu	cct Pro	ctt Leu	Gly 99c	atc Ile 310	caa Gln	gat Asp	agg Arg	cca Pro	ago Ser 315	tta Leu	Gly gga	aac Asn	cct Pro	aaa Lys 320	960
	aga	gaa	gct	tac	gça	aca	att	ctt	çac	tca	gct	cac	gtt	tac	gtċ	tgc	1008

Arg Glu Ala	Tyr Ala Thr	Ile Leu His	Ser Ala	His Val	Tyr Val Cys
	325	•	330	• .	325

	225	
	325 330 335 D ggt gca atc gcc gcg	
•	Sly Ala Ile Ala Ala Ala Gln Ser Ile Arg Gln Ser Gly Ser Thr Arg 340 345 350	1056
10	gac ctt gtt atc ctt gtt gat gac aac atc agc ggt tac cac cgg agt Asp Leu Val Ile Leu Val Asp Asp Asn Ile Ser Gly Tyr His Arg Ser 355 360 365	1104
15	gga cta gaa gcc gcg ggt tgg caa atc cgg acg ata cag agg att cga Gly Leu Glu Ala Ala Gly Trp Glu Ile Arg Thr Ile Glu Arg Ile Arg 370 380	1152
20	aac cct aag gca gag aaa gat gct tac aac gaa tgg aac tac agc aag Asn Pro Lys Ala Glu Lys Asp Ala Tyr Asn Glu Trp Asn Tyr Ser Lys 385 390 395	1200
	ttc cgg cta tgg cag ctg act gat tac gac asa atc att ttc atc gac Phe Arg Leu Trp Gln Leu Thr Asp Tyr Asp Lys Ile Ile Phe Ile Asp . 405 410 415	1248
25	gcg gat ctc tta atc ttg aga aac atc gat ttc ttg ttc tcg atg cct Ala Asp Leu Leu Ile Leu Arg Asn Ile Asp Phe Leu Phe ser Met Pro 420 425 430,	1296
30	gag atc tca gct aca gga aac aat gga act ctg ttt aat tca gga gtt Glu Ile Ser Ala Thr Gly Asn Asn Gly Thr Leu Phe Asn Ser Gly Val 435 / 445	1344
35	atg gtg atc gag cct tgc aac tgt acg ttt cag ctt ctg atg gaa cat Met Val Ile Glu Pro Cys Asn Cys Thr Phe Gln Leu Met Glu His 450 455	1392
40	465 470 475 480	1440
	485 490 Lys His Met Asn Phe Leu	1488
45	aag cat tit tigg att ggc gat gaa gat gac gcg aaa cgc aag aaa aca Lys His Phe Trp Ile Gly Asp Glu Asp Asp Ala Lys Arg Lys Lys Thr 500 505 510	1536
50	gag ctt ttt gga gca gag cct cct gtt ctt tat gtt ctt cat tac ctt 1 Glu Leu Phe Gly Ala Glu Pro Pro Val Leu Tyr Val Leu His Tyr Leu 515 520 525	1584
55	ggg atg aag ccg tgg tta tgt tac cgt gac tac gac tgt aac ttc aac Gly Met Lys Pro Trp Leu Cys Tyr Arg Asp Tyr Asp Cys Asn Phe Asn 530 535	L632
60	tcc gac ata ttc gtt gag ttt gct acc gat atc gct cat cga aaa tgg Ser Asp Ile Phe Val Glu Phe Ala Thr Asp Ile Ala His Arg Lys Trp 545 550 555 560	1680
	tgg atg gtc cac gac gcc atg cca cag gaa-ctt cac-caa ttc tgt tac ITrp Met Val His Asp Ala Met Pro Gln Glu Leu His Gln Phe Cys Tyr	.728

														•			
					565					570					575		
5	ttg Leu	aga	tec Ser	aag Lys 580	caa Ģln	aag Lys	gca Ala	cag Gln	ctg Leu 585	gaa Glu	tat Tyr	gat Asp	Arg Cgc	Arg 590	caa Gln	gca Ala	1776
40	gag Glu	gcc Ala	gca Ala 595	aat Asn	tat Tyr	gcc Ala	gac	ggt Gly 600	cat His	tgg Trp	aaa Lys	ata Ile	aga Arg 605	gta Val	aag Lys	yab gac	1824
10	eeg Pro	aga Arg 610	ttc Phe	aaa Lys	att Ile	tgc Cys	atc Ile 615	gac	aaa Lys	tta Leu	tgt Cys	aat Asn 620	tgg Trp	aaa Lys	agt Ser	atg Met	1872
15	ctg Leu 625	cgg Arg	cat His	tgg Trp	ggc ggc	gaa Glu 630	tca Ser	aat Asn	tgg Trp	act Thr	gac Asp 635	tac Tyr	gag Glu	tct Ser	ttt Phe	gtt Val 640	1920
20	ccc Pro	acc Thr	cca Pro	cca Pro	gcc Ala 645	att Ile	acc Thr	gta Val	gac Asp	egg Arg 650	aga Arg	tca Ser	tca Ser	ctt Leu	ecc Pro 655	GJÀ âāc	1968
25		aac Asn		tga 660	¢gaa	aataa	att a	itaca	atacı	t at	taat	ggat:	: ttc	atga	agtt		2020
	tttt	ggtt	tg a	atto	gttga	t go	gaga	ıttaş	gt	gaata	itca	gtt	jtgta	ac t	atat	:ctt t	2080
	teet	ataç	jtt I	gtto	saat	t ga	ataa	aaaca	a tti	ttt	gca	gttt	aacc	ac		,	2130
30														1.			
35	<212 <212)> 14 L> 6! 2> P! B> A)	59 RT	dopsi	is tl	palia	ana					-					• .
	<400)> 14	1		77	73.0	7.7.0	D TO	בומ	Pro	ሞ ክተ	Thr	Thr	Thr	Glv	Glv	
40	1				5		•			10					15		
40				20					25	Ile				30			
	_		35					40		Gly			45				
45	Pro	Phe 50	Asn	Ile	Ile	Asn	Phe 55	Ser	Thr	Gln	Asp	Fys Lys	Asn	Ser	Ser	Cys	
-10	•	Сув	Phe	Thr	Lys	Phe 70		Ile	Val	Lys	Leu 75	Leu	Leu	Phe	Ile	eo Leu	
	65 Leu	Ser	Ala	Thr		Phe	Thr	Ile	Ile	Tyr		Pro	Glu	Ala	Tyr 95	His	
50	His	Ser	·Leu			Ser	Ser	Ser		90 Trp	Ile	Trp	Arg	Arg 110	Gln	Asp	
	Pro	Arg	Tyr	100 Phe	Ser	Asp	Leu	Asp	105 Ile	Asn	Trp	Asp	Asp			Lys	•
	Thr	Leu	115 Glu	Asp	Ile	Glu	.Glu	Gly Gly	Arg	Thr	Ile	Gly	125 Val	Leu	Asn	Phe	
55	yab	130 Ser	Asn	Glu	. Ile	Gln	135 Arg	Trp	Arg	Glu	Val	140 Ser		Ser	Lув	Asp	
	145					150					155					160 Asp	
60					165	i				170				•	エノコ	Glu	•
ŲŪ				120	•				185	i				Tan	l	Val	
	GID	· GT0		G10	- A C214	. 44.4		1·						_	•		

										29	,					
			19	5				20	0				20	5	••	
			_					3				22/	ı Pri	o Cy		3 Lys
5		_				26.31	,				772	His	Let			ı Ala
	Ala	a Al	a Th	r Vai	l Ala 245	Ala	Se:	r Alá	a Lys	Gly	/ Phe	Phe	Arg	g Gly	, His	240 Val
	Ph	e Ph	e Val	L <i>S</i> ex	r Arg	Cys	Phe	Pro	Ile	250 Pro) Asp	Lev	Phe	Arg	255 Cyt	Lys
10	Ası) Le	u Val 275	l Sez		Arg	Gl	/ Asp	265 [Va]	r L	Leu	Тух	Lys	270) Asr	Leu
			r Lev	•			Lev	. Gln				Gly	285 Ser	•		Leu
15		Lei	-		G1y	Ile	Gli	,			Sex	300 Leu				
					Ala					Ser	315 Ala					
				Ala	Ala				Ile	330 Arg						
20			val	Ile	Leu			Asp	345 Asn					250		_
		Ler	Glu		Ala		Trp	360					366			
25	Asn	Pro			Glu		375					300				_
25					Gln	220					395					4-4
					405 Ile					410					A 7 F	
30				32V	Thr				425		•			420		
			455		Pro			440					445		_	
		420			Glu		455					ACO				
35	~~~				Trp	+± /∪					475					400
					485 Ile					490					4 DE	
40				3 00					505					510		
			DYD		Ala			520					525			
•		230			Trp :		535					54 N				
45	747				Val	330					555					E 6 0
					Asp 2 565					570					E76	
50				200	Gln :				585					500		
30			222		Tyr 1			6UU				•	505			
		970			Ile (3 1 5				1	620				
55	625					530					635					540
				Pro ;	Ala : 645	lle :	thr '	Val 2	Asp .	Arg ; 650	Arg !	er 8	Ser :		Pro (655	Gly
	His	Asn	Leu													
60											•					

<211> 15294

```
<212> DNA
     <213> Artificial Sequence
     <220>
     <220>
     <223> Description of Artificial Sequence: vector
10
     <400> 15
     ggccgggagg gttcgagaag ggggggcacc cccttcggc gtgcgcggtc acgcgcacag 60
     ggcgcagccc tggttaaaaa caaggtttat aaatattggt ttaaaagcag gttaaaagac 120
     aggitagegg iggeegaaaa aegggeggaa aeeetigeaa aigeiggati tietgeeigt 180
     ggacagecce teaaatgtea ataggtgege ceetcatetg teageactet geccetcaag 240
15
     tgtcaaggat cgcgcccctc atctgtcagt agtcgcgccc ctcaagtgtc aataccgcag 300
     ggeacttate eccaggetty tecacateat etytyggaaa etegegtaaa ateaggegtt 360
     ttegeegatt tgegaggetg geeageteea egtegeegge egaaategag eetgeecete 420
     atetgteaac geogegeegg gtgagtegge ceetcaagtg teaacgteeg ceetcatet 480 gteagtgagg geoaagtttt cegegaggta tecacaacge eggeggeege ggtgtetege 540
20
     acacggette gaeggegttt etggegegtt tgeagggeea tagaeggeeg ecageecage 600
     ggcgagggca accagcccgg tgagcgtcgc aaaggcgctc ggtcttgcct tgctcgtcgg 660
     tgatgtactt caccagetee gegaagtege tettettgat ggagegeatg gggaegtget 720
     tggcaatcac gcgcaccccc cggccgtttt agcggctaaa aaagtcatgg ctctgccctc 780
     gggcggacca cgcccatcat gaccttgcca agctcgtcct gcttctcttc gatcttcgcc 840
25
     agcagggega ggatcgtggc atcaccgaac cgcgccgtgc gegggtcgtc ggtgagccag 900
     agtttcagca ggccgccag geggcccagg tcgccattga tgcgggccag ctcgcggacg 960
     tgetcatagt ccacgaegce egtgattttg tagecetgge egaeggeeag eaggtaggee 1020
     qacaggetea tgeeggeege egeegeettt teeteaateg etettegtte gtetggaagg 1080
     cagtacacet tgataggtgg getgeeette etggttgget tggttteate agecateege 1140
30
     ttgccctcat ctgttacgcc ggcggtagcc ggccagcctc gcagagcagg attcccgttg 1200
     agcaccgcca ggtgcgaata agggacagtg aagaaggaac acccgctcgc gggtgggcct 1260
     acticaceta tectgecegg etgacgeegt tggatacace aaggaaagte tacacgaace 1320
     ctttggcaaa atcctgtata tcgtgcgaaa aaggatggat ataccgaaaa aatcgctata 1380
     atgaccecga ageagggtta tgcageggaa aagegeeacg etteecgaág ggagaaagge 1440
35
     ggaCaggtat ccggtaagcg gCagggtCgg aacaggagag cgcacgaggg agcttccagg 1500
     gggaaacgcc tggtatettt atagteetgt egggtttege eacetetgae ttgagegteg 1560
     attitigiga tgetegicag gggggeggag eetatggaaa aaegeeagea aegeggeett 1620
     tttacggttc ctggcctttt gctggccttt tgctcacatg ttctttcctg cgttatcccc 1680
     tgattetgtg gataaccgta ttaccgcctt tgagtgagct gataccgctc gccgcagccg 1740 aacgaccgag cgcagcgagt cagtgagcga ggaagcggaa gagcgccaga aggccgccag 1800
40
     agaggccgag cgcggccgtg aggcttggac gctagggcag ggcatgaaaa agcccgtagc 1860
     gggetgetae gggegtetga egeggtggaa agggggaggg gatgttgtet acatggetet 1920
     gctgtagtga gtgggttgcg Ctccggcagc ggtcctgatc aatcgtcacc ctttctcggt 1980
     ccttcaacgt tcctgacaac gagcctcctt ttcgccaatc catcgacaat caccgcgagt 2040
45
     ccctgctcga acgctgcgtc cggaccggct tcgtcgaagg cgtctatcgc ggcccgcaac 2100
     ageggegaga geggageetg tteaacggtg eegeegeget egeeggeate getgtegeeg 2160
     gootgotoot caagcacggo cocaacagtg aagtagotga tigtoatcag cgcattgacg 2220
     gcgtccccgg ccgaaaaacc cgcctcgcag aggaagcgaa gctgcgcgtc ggccgtttcc 2280
     atotgoggtg egeceggteg egtgeeggea tggatgegeg egecategeg gtaggegage 2340
50
     agegeetgee tgaagetgeg ggeatteeeg atcagaaatg agegeeagte gtegtegget 2400
     ctcggcaccg aatgcgtatg attctccgcc agcatggctt cggccagtgc gtcgagcagc 2460
     geocgettgt teetgaagtg coagtaaage geoggetget gaaececeaa cegtteegee 2520
     agtttgcgtg tcgtcagacc gtctacgocg acetcgttca acaggtccag ggcggcacgg 2580
     atcactgtat toggotgcaa cittgtcatg cittgacactt tatcactgat aaacataata 2640
55
     tgtccaccaa Cttatcagtg ataaagaatc cgcgcgttca atcggaccag cggaggctgg 2700
     toeggagged agaegtgaaa eccaacatac coetgategt aattetgage actgtegege 2760
    togacgotgt cggcatcggc otgattatgc cggtgctgcc gggcctcctg cgcgatctgg 2820
     ttcactcgaa cgacgtcacc gcccactatg gcattetgct ggcgctgtat gcgttggtgc 2880
     aatttgcctg cgcacctgtg ctgggcgcgc tgtcggatcg tttcgggcgg cggccaatct 2940
60
     tgetegtete getggeegge gecaagatet ggggaaceet gtggttggea tgeacataca 3000
     aatggacgaa cggataaacc ttttcacgcc cttttaaata tccgattatt ctaataaacg 3060
     ctettttete traggittae cegecaatat ateetgicaa acactgatag titaaactga 3120
```

	agacagaaa	- casastat					
	accostosco	. cgacaaccc	accargage	y gagaattaag	ggagtcacgt	: tatgacccc	3180
	300300300	r caaaacaaac	egecetacgt	ttggaactga	ggagtcacgt	acgttqaaqq	3240
	ttatactact	, cegacergae	i ttcccgatct	agtaacatag	atgacacege	gegegataat	3300
[.] 5	estatosas	regegegete	tattttgttt	: tctatcgcgt	atgacaccgo attaaatgta	taattocoo	3360
3	tostenaces	taaaaaccca	i teteataaat	aacgtcatgo	actaeatgta attacatgtt	aattattaca	3420
	tgettaaegt	aartcaacag	, asattatatg	r ataatcatcg	caagaccggc	Bacaccatta	3490
	aatcttaaga	. aactttatto	ccaaatgttt	gaacgatcgo	ggaaattcga	acteggatte	3540
	atcatgttac	aaacttttt	gctgtgagca	gtagatatoo	ggaaarrcga aaacccggag	doceggeace	3540
40	atctgataga	taaagcacat	agcttgggtt	tacacette	ggtggatgta	gacctaaagt	3600.
10	atgcaagcaa	taatgccact	gatggcctca	atogetros	tattggccaa	grrcacagte	3660
	aatectactt	tcatgctgga	gagcaagggt	accateeatt	gtgggatage	ggtteteaag	3720
	actatgccaa	ttgggaggtt	ctteatttee	ttatttage	grgggatage	aggetgttca	3780
	agtataactt	tgacggattt	castttrata	Gastosett-	tatgctgtat	tggctagaag	3840
	gaatcaatat	gggatttaca	Transchate	gaacaactt	targergrat	gttcatcatg	3900
15	atoctotoot	ctatttaata	tteresee	argagtattt	cagegagget	acagatgttg	3960
20							
20							
25							
	atgatgagaa	gaaggttgtt	atatttaaac	CTCCtcggaaa	ggtatttgta	agcagcatgg	4620
	acccaaataa	Cacatacgaa	gagteteaaa	grage dates	ggcacccgca	ttcaacttcc	4680
	ttgcactgga	cagtgatget	toorsates	rrddardrda	cttgccaggg	aagtacagag	4740
30							
25							
35			4 CHANGE CCC	CUCCCCCTTTA	277277744		
		3-3-3-4-4-4	44 LAGLU LAL	acceaceaca	COTTO PARAME		
	J J J		-446	aci.uardeda	27777744		
				CHUMEONARE	T 3 7 7 3 7 5 8 8 8 8	~~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~	
40 ⁻							
	tgcatatggc	aattgctgat	aaatggattg	anttactors	gaaacgggat	gactategge	5520
	gagtgggtga	tattottcat	acactgaces	atacaacaa	agarcadast.	gaggartgga	5580
	acoctoaaao	tcatgatcaa	oct ct act ca	acagaagarg	gtoggaaaag	rgrgrrrcat	5640
	acaaogatat	gratgatest	stantates	gryaraaaac	tatagcattc	tggctgatgg	5700
45		3 40043 70 70 70	acuncicio	MCADACCETC.	22724~~~		
		3	uccauaccca	CHACTATOOK	2222444		
		3334446	CLUMMUCALL	CEGRATAGE	アクラナナナハススト		
		-34-33-4-4	Aragrececo.	gaaaccaarr	Canttabase	~~~*~~~~	F040
	22-2-6-20	A CORRECTOR	ucauaacac e	CARGATACCA	TAAA## ~~~~		
EΛ	222craraca	gracectogaa	dataaatato	agtttatgar -	ttdadaaaaa	~~~ ~	~^~
50		ころにひろるに	#99acqactq	tatttoaaaa	accessors .	~	C 7 C A
	CCCCCCC	ccyyacaaaa	agctattcar	actategest	agactacata .	33 <i>6</i> 004666	6100
	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	-209986	LCAGALGALC	Cactttttdd	tancetonno.	~~~~++~~+~	C 2 4 A
	araacyccya	acallicate	tttgaaqqat	qqtatqatqa	teatectear ·	tasstratee	6300
	tgtatgcacc	tagtagaaca	gcagtggtct	atocactaot	202000000	ccaarracyy **	6360
<b>55</b>	aagaagaagt	agcagtagta	gaagaagtag	tagtagaage	-arrandada ;	anappet = - t	03 <b>0</b> U
	cgcgttgaaa	gatttgaagg	Ctacatacat	Ctadacter-	waaraaraacg	aaccegtgat	042U·
	attggaggag	atgagtasss	attaccanet	aaaaaccaa	cetgeatgaa :	accagaaata	6480
	attggaggag	3~3~44AA	arrancacte	yergagetgt	grgagrgagt	gagtgagaat	6540
	gaggaggtgc	ttaccence-	cycaycaggt	ccagtgaca	cgtgtcaaga	gaatagcggg	6600
60	cygocacecc	ccaycayaag	gcaactgtgg	acactotatt	atagggaaat i	octcatcoso	EEEN
.00	agrarrary	gecoure	LYCCGACTCA	Caactaaaat	teaactteer /	anttonionto.	572A
	ggcccgcccg	gttelgtete	ccagtateta	aaaaactaaa -	<b>ぐぐるみのたのかのた…</b> 。	cat sacrata	C-7.0 A.
	ccacttgaca	trectatgte	tcgtgttaat	taaattatta	ttatagtaat	taaaaataat	6840
					_		

atctaggtac tggtactggt occtocotoc actagaatat tagttacttc coccttagct 6900 ttgtattcca aattactgta aatatatttt ctaatttttt acgacaaaca agatctaatt 6960 atgaatgcac aattetaaag gttgaataca ttactttact tggtttagcc tatattaagt 7020 tgcattttag tattaagatt gagatgcatg gttctattac aaaattgata cactgctaaa 7080 ggaaggatgg ttaaamacaa cattcaatgt ttgttacatt tcttcctatt gtatttttt 7140 5 tttaacgago ttoccetata catcataaca tetotocett coacttegca egaaaaaaaa 7200 atacccaaac aggaagatac tgtcaagtat atccatagat gaggacttaa tggataggct 7260 tttcqaggat tcataaatca taatatctgg cggaggagtc aattaaatac ttgtggtttg 7320 tatoctgatt actcogtcaa cagocaaata gaaaagtttg aaaagagaga aaggatttgg 7380 10 tacaagatac tgttgcattt gttaagtaat gaacaaaacg gagtaacata attttctatc 7440 tegttaaage tteaegetge egeaageaet eagggegeaa gggetgetaa ggaageggaa 7500 cacgtagaaa gccagtccgc agaaacggtg ctgaccccgg atgaatgtca gctactgggc 7560 tatetggaca agggaaaacg caagegeaaa gagaaageag gtagettgea gtgggettae 7620 atggcgatag ctagactggg cggttttatg gacagcaagc gaaccggaat tgccagctgg 7680 ggcgccetct ggtaaggttg ggaagccctg caaagtaaac tggatggett tcttgccgcc 7740 15 aaggatotga tggogcaggg gatcaagato atgagoggag aattaaggga gtcacgttat 7800 gacccccgcc gatgacgcgg gacaagccgt tttacgtttg gaactgacag aaccgcaacg 7860 ttgaaggage cactcagecg egggtttetg gagtttaatg agetaageae ataegtcaga 7920 aaccattatt gcgcgttcaa aagtcgccta aggtcactat cagctagcaa atatttcttg 7980 20 tcaaaaatgc tccactgacg ttccataaat tcccctcggt atccaattag agtctcatat 8040 tcactotcaa tccagatoto gactotagto gagggcccat gggagottgg attgaacaag 8100 atggattgca egeaggttet eeggeegett gggtggagag getattegge tatgaetggg 8160 cacaacagac aateggetge tetgatgeeg cegtgtteeg getgteageg caggggegee 8220 eggttetttt tgtcaagace gacetgteeg gtgcoetgaa tgaactgeag gacgaggeag 8280 25 egeggetate giggetggee acgaeggges treettgege agetgtgete gaegttgtea 8340 ctgaagcggg aagggactgg ctgctattgg gcgaagtgcc ggggcaggat ctcctgtcat 8400 ctcaccttgc tcctgccgag aaagtatcca tcatggctga tgcaatgcgg cggctgcata 8460 egettgatee ggetacetge ecattegace accaagegaa acategeate gagegageae 8520 gtactcggat ggaagccggt cttgtcgatc aggatgatct ggacgaagag'catcaggggc 8580 30 tegegocage egaactgite gecaggetea aggegocat gecegaegge gaggateteg 8640 tegtgaceca tggcgatgcc tgcttgccga atatcatggt ggaaaatggc cgcttttctg 8700 gattcatcga ctgtggccgg ctgggtgtgg cggaccgcta tcaggacata gcgttggcta 8760 cccqtgatat tgctgaagag cttggcggcg aatgggctga ccgcttcctc gtgctttacg 8820 gtategoege tecegatteg cagegoateg cettetateg cettettgae gagttettet 8880 gagegggace caagetaget tegaeggate eccegatgag etaagetage tatateatea 8940 35 atttatgtat tacacataat ategeactca gtettteate taeggeaatg taccagetga 9000 tataatcagt tattgaasta tttctgaatt taaacttgca tcaataaatt tatgtttttg 9060 aagatgggaa ttaattcact ggccgtcgtt ttacaacgtc gtgactggga aaaccctggc 9180 gttacccaac ttaatcgcct tgcagcacat coccctttog ccagctggcg taatagcgaa 9240 40 gaggeeegea eegategeee tteecaacag ttgegeagee tgaatggege eegeteettt 9300 cgctttette ectteette tegecaegtt egeeggettt eccegteaag etetaaateg 9360 ggggeteect ttagggttee gatttagtge tttaeggeae etegaeecea aaaaaettga 9420 tttgggtgat ggttcacgta gtgggccatc gccctgatag acggtttttc gccctttgac 9480 gttggagtcc acgttcttta atagtggact cttgttccaa actggaacaa cactcaaccc 9540 45 tatctoggge tattetettg atttataagg gattttgeeg attteggaac caccateaaa 9600 caggattttc gcctgctggg gcaaaccagc gtggaccgct tgctgcaact ctctcagggc 9660 caggeggtga agggeaatea getgttgeee gtetcaetgg tgaaaagaaa aaceacocca 9720 gtacattaaa aacgtccgca atgtgttatt aagttgtcta agcgtcaatt tgtttacacc 9780 acaatatate etgecaccag ecagecaaca geteceegae eggeageteg geacaaaate 9840 50 accactcgat acaggcagcc catcagtccg ggacggcgtc agcgggagag ccgttgtaag 9900 qcggcagact ttgctcatgt taccgatgct attcggaaga acggcaacta agctgccggg 9960 tttgaaacac ggatgatete geggagggta geatgttgat tgtaacgatg acagagegtt 10020 getgeetgtg atcasatate atcteeteg cagagatecg aattateage ettettatte 10080 atttctcgct taaccgtgac aggctgtcga tcttgagaac tatgocgaca taataggaaa 10140 55 tegetggata aageegetga ggaagetgag tggegetatt tetttagaag tgaaegttga 10200 cgatatcaac teccetatec attgeteace gaatggtaca ggteggggac cegaagttee 10260 gactgtcggc ctgatgcate cccggctgat cgaccccaga tctagatctg gggctgagaa 10320 ageccagtaa ggaaacaact gtaggttega gtegegagat eeeceggaac caaaggaagt 10380 aggttaaacc cgctccgatc aggccgagcc acgccaggcc gagaacattg gttcctgtag 10440 60 gcatcgggat tggcggatca aacactaaag ctactggaac gagcagaagt cctccggccg 10500 ccagttgcca ggcggtaaag gtgagcagag gcacgggagg ttgccacttg cgggtcagca 10560

				44			
	eggtteega	a coccatoga	a accgcccc	g ccaggcccg	c tacaacaca	g acaggateta	30600
	acaccacac	t tggtgtcaa	c accaacagc	g ccacgcccg	c agtteegea	g acaggateta a atagececca	10000
	ggaccgcca	t caatcgtat	c gggctacct	a gcagagcgg	c agagatgaa	atagococca acgaccatca	10740
5	geggetgea	c agcgcctac	a afcaccaca	a ccccgcccg	g Caggoggtad	o acgaccatca g accgaaataa	10740
•	acaecaayc	t ccagaatag	c gaaatatta	a gtgcgccga	gatgaagat	y accgaaataa y cgcatccacc	10860
	agaceeegg	L Lygaatetg	cggacgatca	a tcacgagça:	a taaacccgc	ggcaacgccc ggcaacgccc	10920
	gcagcagca	c accggcgace	c cctcggcct	gctgttcgg;	Ctccacgaa	acgccggaca g acgccggaca	10020
	tatagagaga	c grgagcgtc	ttggggccg	cctcctgtt	gaagaccga	acgooggaca agoocaatga	11040
10	adat accept	e daracadaco	g ccgaatgcca	a cggcatctc	7 Caaccgttce	agcccaatga gcgaacgcct	11100
	Coargage	r rrrerected	, tgctcgtaaa	ı cggacccgaa	a catctctgge	gcgaacgcct gctttcttca	11150
	gggeegaea	coggatoto	y cggaaatcct	: gcacgtcgg(	cgctccaagc	getttettea egtegaatet	11220
	Catacataa	- cacaatege	aattttaat	ctctgtttat	cggcagttcg	cgtcgaatct tagagege	11280
`	2225090000	gagegataet	gagcgaagca	agtgcgtcga	gcagtgcccg	tagagegege cttgtteetg	11340
15	gratttaca	aaaycyctgg	ctgctgaac	cccagccgg	actgacccca	caaggcccta	11400
	Teacher teacher	t tgcaccaggt	catcattgad	ccaggcgtgt	tccaccagge	caaggcccta cgctgcctcg	11460
	teeesante	eaggettege	cgacctgctc	gcgccactto	ttcacgcggg	cgctgcctcg tggaatccga	11520
	atagtgata	aggeggaagg	tttccagett	gagcgggtac	ggeteeeggt	gcgagctgaa	11580
	cotetache	accegeeggg	. ccarcaaca	cagettgegg	tacttctccc	gcgagctgaa atatgaattt	11640
20	CGCGCCGG	regecageaa	acagcacgac	gatttcctcg	togatoagga	atatgaattt cctggcaacg	11700
20	gracgeetee	rigecacggt	ccaggacgcg	gaagcggtgc	agcagcgaca	cctggcaacg ccgattccag	11760
	graceasts.	caarcaaaca	, tgaagcccat	cgccgtcgcc	tgtaggcgcg	ccgattccag acaggcattc	11820
	Czeggeete	gegeaatace	ggccattgat	cgaccagccc	aggtcctggc	acaggcattc aaagctcgta	11880
	esecuciones esecuciones	acarcades	caccastada	ggtgcgcttc	gcgtactcca	acacctgctg	11940
25	Cttattaaa	tegreategt	cggcccgcag	ctcgacgccg	gtgtaggtga	acacctgctg tcttcacgtc	12000
	cetestegacy	cyyaaaacga	ccttgttttg	cagcgcctcg	cgcgggattt	tetteacete	12060
	CCCCCCCCCCC	ayyycagagc	aaaccararc	gtttggcatc	gctcgcatcg	tettetteeg tetceggeea	12120
	caacacaca	rcasscaage.	. aaagctgcat	ttccttgatc	tgctgcttcg	tgtccggcca tgtgtttcag	12180
	Cttcttcttc	rgertggeet	cgctgacctg	ttttgccagg	teetegeegg	tgtgtttcag cggtttttcg	12240
30	ctcctattca	accaragece	cccgcgtgtc	gatggtcatc	gacttegeea	aacctgccgc	12300
	agccagtter	agacyacyeg	aacgeteeac	aacaaccaar	ggcgcgggca	gggcaggggg aacctgccgc	12360
35							
40							
. •							
45							
50							
			Lactatic	CCCCCCCCC	<u> </u>	<b>_</b>	
			awacuaacii:	CAMETERATA	**********		
55			CCCCUGGGG	ULCCCCAPAN .	TTARAMENES.		~
	J			<u>uuluaraeta</u>	~~~~~~~~ .		
				LLCLGEBRA	2TC2CCFCtc .		^^~
		2222222444	M COOLAK COO	aaucaccacc	MM300800		
		US COMMUNICAL	44Caaccuca	GCCCACTTAC .	ひとだけのたたったっ ・	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	4040
60							
	· · · · · - J J J - J J	Dagger and Ca.	vgaccucaty .	tcacttaaaa	AACTCACCCC (	resetences 1	47 CO
		eremanaging .	Caytacate	aecorcraca	CCCDDDDRAAA	4 <b>1</b>	4556
	JJ300g	יייייייייייייייייייייייייייייייייייייי	acaccadede	-Adccdrrrr	acgcgtatga (	cagtctccgg 1	4280
							•

	aagacggttg	ttgcgcacgt	atteggtgaa	cgcactatgg	cgacgctggg	gcgtcttatg	14340
•	agcctgctgt	caccctttga	cgtggtgata	tggatgacgg	atggctggcc	gctgtatgaa	14400
						aattgagcgg	
_						gtcgttctca	
5	aaatcggtgg	agctgcatga	caaagtcatc	gggcattatc	tgaacataaa	acactatcaa	14580
	taagttggag	tcattaccca	attatgatag	aatttacaag	ctataaggtt	attgtcctgg	14640
	gtttcaagca	ttagtccatg	caagttttta	tgctttgccc	attotataga	tatattgata	14700
	agcgcgctgc	ctatgccttg	ccccctgaaa	tccttacata	cggcgatatc	ttctatataa	14760
						cctcatcctc	
10	ttcatcctct	tcgtcttggt	agctttttaa	atatggcgct	tcatagagta	attetgtasa	14880
	ggtccaattc	tcgttttcat	acctcggtat	aatcttacct	atcacctcaa	atggttcgct	14940
	gggtttatcg	cacccccgaa	cacgagcacg	gcacccgcga	ccactatgcc	aagaatgccc	15000
	aaggtaaaaa	ttgccggccc	cgccatgaag	tccgtgaatg	ccccgacggc	cgaagtgaag	15060
	ggcaggccgc	cacccaggcc	gccgccctca	ctgcccggca	cctggtcgct	gaatgtcgat	15120
15	gċcagcacct						
	gttactgccc	cgatecegge	aatggcaagg	actgccagcg	ctgccatttt	tggggtgagg	15240
	ccgttcgcgg	CCGAGGGGCG	cagcccctqq	ggggatggga	ggcccgcgtt	ageg	15294

Figure 1




Figure 2

į	Mother va-	Dm	Fructose %		Sucrose %
ĺ	riety	%	of	Glucose %	of
Variety/line		}	Ðm	of Dm	Dm
Producent		27,8	1,8	2,2	. 3,1
Prevalent		27,1	1,2	1,7	2,7
Dinamo		27,7	0,8	1,7	2,2
Kuras		27,9	1,8	2,2	2,4
AM98-2012	Producent	22,3	3,1	4,0	4,7
AM98-2019	Prevalent	18,4	2,9	4,4	3,8
AM98-2021	Prevalent	17,1	3,1	5,9	3,5
AM99-2002	Dinamo	19,3	2,4	3,8	3,0
AM99-2003	Dinamo	18,8	2,5	5,5	3,2
AM99-2004	Dinamo	11,7	4,0	6,8	2,1
AM00-2040	Kuras	21,0	5,0	6,3	3,5
AM00-2041	Kuras	19,3	5,6	7,1	3,0

#### Figure 3

3 a) pHS1 for gene-inhibition of StGH1

# pHS1



3 b) pHS2 for gene-inhibition of StGH2

# pHS2



3 c) pHS3 for over-expression of StGH1

10

pHS3



3 d) pHS4 for over-expression of StGH2

# pHS4



Figure 4




Figure 5




Figure 6




Figure 7




Figure 8





Figure 9



pHAbe12A 7756 bp

Enhanced amylose production in plants

#### **Abstract**

The invention relates to methods for increasing the amylose content in plants, preferably in potato plants, by expressing a starch biosynthesis enhancing protein. The invention furthermore relates to an expression cassette expressing the polypeptide in potato plants, preferably in the tubers, the transgenic plants expressing the polypeptide and to the use of said transgenic plants for the production of fine chemicals especially other than native starches.

07-MÄR-2003 16:59

PCT/EP2004/002096

# This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

### **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

D	efects in the images include but are not limited to the items checked:
	☐ BLACK BORDERS
	☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
	☐ FADED TEXT OR DRAWING
	☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
	☐ SKEWED/SLANTED IMAGES
	COLOR OR BLACK AND WHITE PHOTOGRAPHS
	GRAY SCALE DOCUMENTS
	LINES OR MARKS ON ORIGINAL DOCUMENT
	REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

## IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.